

Population health profile of the North East Valley

Division of General Practice: supplement

Population Profile Series: No. 39a

PHIDU

March 2007



PHIDU



Australian Government

Australian Institute of
Health and Welfare



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA

© Commonwealth of Australia 2007

This work may be reproduced and used subject to acknowledgement of the source of any material so reproduced.

National Library of Australia Cataloguing in Publication entry

Population health profile of the North East Valley Division of General Practice: supplement.

Bibliography.

ISBN 9 78073089 6371 (web).

1. Public health - Victoria, Northeastern - Statistics. 2. Health status indicators - Victoria, Northeastern - Statistics. 3. Health service areas - Victoria, Northeastern. 4. Victoria, Northeastern - Statistics, Medical. I. Public Health Information Development Unit (Australia). (Series : Population profile series ; no. 39a).

362.1099455

ISSN 1833-0452 Population Profile Series

Public Health Information Development Unit, The University of Adelaide
A Collaborating Unit of the Australian Institute of Health and Welfare

This profile was produced by PHIDU, the Public Health Information Development Unit at The University of Adelaide, South Australia. The work was funded under a grant from the Australian Government Department of Health and Ageing. The views expressed in this profile are solely those of the authors and should not be attributed to the Department of Health and Ageing or the Minister for Health and Ageing.

Interpretation of differences between data in this profile and similar data from other sources needs to be undertaken with care, as such differences may be due to the use of different methodology to produce the data.

Suggested citation:

PHIDU. (2007) *Population health profile of the North East Valley Division of General Practice: supplement*. Population Profile Series: No. 39a. Public Health Information Development Unit (PHIDU), Adelaide.

Enquiries about or comments on this publication should be addressed to:

PHIDU, The University of Adelaide, South Australia 5005
Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au

This publication, the maps and supporting data, together with other publications on population health, are available from the PHIDU website (www.publichealth.gov.au).

Published by Public Health Information Development Unit, The University of Adelaide

Contributors: Anthea Page, Sarah Ambrose, Kristin Leahy and John Glover

Population health profile of the North East Valley Division of General Practice: supplement

This profile is a supplement to the *Population health profile of the North East Valley Division of General Practice*, dated November 2005, available from www.publichealth.gov.au. This supplement includes an update of the population of the North East Valley Division of General Practice, as well as additional indicators and aspects of the Division's socioeconomic status, use of GP services and health. The contents are:


- Population [updated to June 2005]
- Additional socio-demographic indicators
- Unreferred attendances – patient flow/ GP catchment
- Additional prevalence estimates: chronic diseases and risk factors combined
- Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions
- Avoidable mortality

For further information on the way Division totals in this report have been estimated, please refer to the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Population

The North East Valley Division had an Estimated Resident Population of 233,270 at 30 June 2005.

Figure 1: Annual population change, North East Valley DGP, Melbourne, Victoria and Australia, 1991 to 1996, 1996 to 2001 and 2001 to 2005




Over the five years from 1991 to 1996, the Division's population increased by 0.2% on average each year, well below the levels in Melbourne (0.8%), Victoria (0.6%), and Australia as a whole (1.2%). From 1996 to 2001, the annual percentage increase was 0.5%, compared to increases of 1.3% for Melbourne, and 1.2% for Victoria. From 2001 to 2005 the Division's population decreased by 0.2%, compared to annual increases of 1.0% for Melbourne and Victoria, and 1.1% for Australia.

Table 1: Population by age, North East Valley DGP and Australia, 2005

Age group (years)	North East Valley DGP		Australia	
	No.	%	No.	%
0-14	42,136	18.1	3,978,221	19.6
15-24	32,264	13.8	2,819,834	13.9
25-44	70,829	30.4	5,878,107	28.9
45-64	58,520	25.1	4,984,446	24.5
65-74	15,118	6.5	1,398,831	6.9
75-84	10,777	4.6	954,143	4.7
85+	3,625	1.6	315,027	1.5
Total	233,270	100.0	20,328,609	100.0

As shown in the accompanying table and the age-sex pyramid (Figure 2), the North East Valley DGP had slightly fewer children at ages 0 to 14 years (18.1%) compared to Australia as a whole (19.6%) (Table 1). Conversely, the proportion of the Division's population aged 25 to 44 years (30.4%) was slightly higher than for Australia (with 28.9%). The proportions in other age groups were consistent with those for Australia.


Figure 2: Population in North East Valley DGP and Australia, by age and sex, 2005



The age distribution of the Division's population is very similar to that for Australia overall. The only notable differences are:

- at younger ages – lower proportions of children and young people aged 0 to 19 years; and
- from 20 to 54 years – slightly higher proportions of both males and females.

Figure 3: Population projections for North East Valley DGP, by age and sex, 2005 and 2020




The population projections for the Division show a number of changes in age distribution, with the 2020 population projected to have:

- at younger ages – lower proportions of children, young people, and young adults aged 0 to 24 years;
- from 30 to 54 years – lower proportions of males and females; and
- at older ages – higher proportions of both males and females aged 60 years and over, with the largest differentials at ages 60 to 74 years.

Additional socio-demographic indicators

Please refer to the earlier *Population health profile of the North East Valley Division of General Practice*, dated November 2005, available from www.publichealth.gov.au, for other socio-demographic indicators.

Figure 4: Index of Relative Socio-Economic Disadvantage, North East Valley DGP, 2001



One of four socioeconomic indexes for areas produced at the 2001 ABS Census is the Index of Relative Socio-Economic Disadvantage.

The North East Valley DGP has an index score of 1062, above the score for Australia of 1000: this score varies across the Division, from 953 in the most disadvantaged areas to 1131 in the least disadvantaged areas.


Note: each 'quintile' comprises approximately 20% of the population of the Division.

A new indicator, produced for the first time at the 2001 ABS Census, shows the number of jobless families with children under 15 years of age. There were markedly fewer jobless families in the North East Valley DGP (10.6%), compared to Melbourne as a whole (14.7%) (Figure 5, Table 2).

With the introduction of the 30% rebate for private health insurance premiums, there was a once-off registration process, providing information of the postcode and residence of those who had such insurance (these data are not available at this area level for later dates). In 2001, the Division had a notably higher proportion of people with private health insurance (57.6%), compared to Melbourne (49.2%) (Figure 5, Table 2).

Figure 5: Socio-demographic indicators, North East Valley DGP, Melbourne, Victoria and Australia, 2001

Jobless families with children under 15 years old



Private health insurance, 30 June





Table 2: Socio-demographic indicators, North East Valley DGP, Melbourne, Victoria and Australia, 2001


Indicator	North East Valley DGP		Melbourne		Victoria		Australia	
	No.	%	No.	%	No.	%	No.	%
Jobless families with children under 15 years old	2,574	10.6	52,418	14.7	77,142	15.4	357,563	17.4
Private health insurance (30 June)	128,722	57.6	1,653,598	49.2	2,196,890	47.5	8,671,106	46.0

Details of the distribution of jobless families and of the population covered by private health insurance are shown by Statistical Local Area (SLA) in Maps 1 and 2, respectively.

Map 1: Jobless families with children under 15 years of age by SLA, North East Valley DGP, 2001



Map 2: People covered by private health insurance by SLA, North East Valley DGP, 30 June 2001



GP services to residents of the North East Valley DGP

The following tables include information, purchased from Medicare Australia, of the movement of patients and GPs between Divisions. Note that the data only include unreferred attendances recorded under Medicare: unreferred attendances not included are those for which the cost is met by the Department of Veterans' Affairs or a compensation scheme; or are provided by salaried medical officers in hospitals, community health services or Aboriginal Medical Services, and which are not billed to Medicare. At any attendance, one or more services may have been provided.

Nearly three quarters (71.5%) of all unreferred attendances to residents of the North East Valley DGP were provided in the Division (ie. by a GP with a provider number in the Division): this represented 825,017 GP unreferred attendances (Table 3). A further 9.9% of unreferred attendances to residents were provided by GPs with a provider number in Northern DGP.

Table 3: Patient flow – People living¹ in North East Valley DGP by Division where attendance occurred², 2003/04

Division		Unreferred attendances	
Number	Name	No.	% ³
302	North East Valley DGP	825,017	71.5
308	Northern DGP	114,460	9.9
301	Melbourne DGP	91,859	8.0
303	Inner Eastern Melbourne DGP	33,862	2.9
310	Whitehorse DGP	23,936	2.1
304	Southcity DGP	14,241	1.2
307	North West Melbourne DGP	11,952	1.0
Other	..	38,154	3.3
Total	..	1,153,481	100.0

¹ Based on address in Medicare records

² Division of GP based on provider number

³ Proportion of all unreferred attendances of patients with an address in Division 302 by Division in which attendance occurred

Almost three quarters (72.4%) of unreferred attendances provided by GPs with a provider number in North East Valley DGP were also to people living in the Division (ie. their Medicare address was in the Division) (Table 4). A further 12.4% of unreferred attendances provided by GPs in the Division were to residents of Northern DGP, with 3.8% to people living in Inner Eastern Melbourne DGP.

Table 4: GP catchment – Unreferred attendances provided by GPs¹ in North East Valley DGP by Division of patient address², 2003/04

Division		Unreferred attendances	
Number	Name	No.	% ³
302	North East Valley DGP	825,017	72.4
308	Northern DGP	141,697	12.4
303	Inner Eastern Melbourne DGP	43,603	3.8
301	Melbourne DGP	30,636	2.7
310	Whitehorse DGP	25,833	2.3
307	North West Melbourne DGP	19,654	1.7
306	Western Melbourne DGP	6,300	0.6
Other	..	46,921	4.0
Total	..	1,139,661	100.0

¹ Division of GP based on provider number

² Based on address in Medicare records

³ Proportion of all unreferred attendances to GPs with a provider number in Division 302 Division of patient address

Additional prevalence estimates: chronic diseases and risk factors combined

Please refer to the earlier *Population health profile of the North East Valley Division of General Practice*, dated November 2005, available from www.publichealth.gov.au, for the separate prevalence estimates of chronic disease; measures of self-reported health and risk factors. The process by which the estimates have been made, and details of their limitations, are also described in the 'Notes on the data' section of this earlier profile.

In this section two estimates, which combine the prevalence of selected chronic diseases with a risk factor, are shown for the Division. The measures are of people who *had asthma and were smokers*, and people who *had type 2 diabetes and were overweight or obese*: note that the estimates have been predicted from self-reported data, and are not based on clinical records or physical measures.

It is estimated that there were relatively fewer people in North East Valley DGP who had asthma and were smokers, compared to Melbourne or Australia as a whole (Figure 6, Table 5): that is, the prevalence rates per 1,000 population were lower. Similarly, there were fewer people in North East Valley DGP who had type 2 diabetes and were overweight/ obese, compared to Melbourne or Australia.

Figure 6: Estimates of selected chronic diseases and risk factors, North East Valley DGP, Melbourne and Australia, 2001



Table 5: Estimates of selected chronic diseases and risk factors, North East Valley DGP, Melbourne, Victoria and Australia, 2001

Variable	North East Valley DGP		Melbourne		Victoria		Australia	
	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ¹
Had asthma & smoked ³	4,196	17.5	66,240	18.4	95,664	19.9	397,734	20.8
Had type 2 diabetes & were overweight/ obese ⁴	3,130	14.6	50,057	15.6	69,192	15.1	283,176	15.2

¹ No. is a weighted estimate of the number of people in North East Valley DGP reporting these chronic conditions/ with these risk factors and is derived from synthetic predictions from the 2001 NHS

² Rate is the indirectly age-standardised rate per 1,000 population

³ Population aged 18 years and over

⁴ Population aged 15 years and over

Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions

The rationale underlying the concept of avoidable hospitalisations is that timely and effective care of certain conditions, delivered in a primary care setting, can reduce the risk of hospitalisation. Admissions to hospital for these ambulatory care sensitive (ACS) conditions can be avoided in three ways. Firstly, for conditions that are usually preventable through immunisation or nutritional intervention, disease can be prevented almost entirely. Secondly, diseases or conditions that can lead to rapid onset problems, such as dehydration and gastroenteritis, can be treated. Thirdly, chronic conditions, such as congestive heart failure, can be managed to prevent or reduce the severity of acute flare-ups to avoid hospitalisation.

This measure does not include other aspects of avoidable morbidity, namely potentially preventable hospitalisations (hospitalisations resulting from diseases preventable through population based health promotion strategies, e.g. alcohol-related conditions; and most cases of lung cancer) and hospitalisations avoidable through injury prevention (e.g. road traffic accidents).

For information on the ambulatory care sensitive conditions and ICD codes included in the analysis in this section, please refer to the *Atlas of Avoidable Hospitalisations in Australia: ambulatory care-sensitive conditions*, available from www.publichealth.gov.au.

In 2001 to 2002, the 5,766 admissions from ambulatory care sensitive (ACS) conditions accounted for 7.5% of all admissions in the North East Valley DGP (Table 6, Figure 7), notably lower than the levels in Victoria (8.8%) and Australia (8.7%).


Table 6: Avoidable¹ and unavoidable hospitalisations, North East Valley DGP, Victoria, and Australia, 2001/02

Category	North East Valley DGP			Victoria			Australia		
	No.	Rate ²	%	No.	Rate ²	%	No.	Rate ²	%
Avoidable ¹	5,766	2,469.2	7.5	145,135	2,983.2	8.8	552,786	2,847.5	8.7
Unavoidable	71,022	29,847.2	92.5	1,510,437	31,088.3	91.2	5,818,199	29,970.7	91.3
Total	76,788	32,322.6	100.0	1,655,572	34,071.5	100.0	6,370,985	32,818.2	100.0

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

Figure 7: Avoidable hospitalisations¹, North East Valley DGP, Victoria and Australia, 2001/02




The rate of avoidable hospitalisations in North East Valley DGP, 2,469.2 admissions per 100,000 population, is markedly lower than the rates for Victoria (a rate of 2,983.2) and notably below that for Australia (2,847.5).

¹ Admissions resulting from ACS conditions

Diabetes complications, chronic obstructive pulmonary disease, dehydration and gastroenteritis, dental conditions and congestive heart failure were the five conditions with the highest rates of avoidable hospitalisations in the North East Valley DGP (Figure 8, Table 7).

Table 7 shows the number, rate and proportion of avoidable hospitalisations, for the individual ACS conditions, as well as the vaccine-preventable; acute; and chronic sub-categories. Two-thirds of avoidable hospitalisations are attributable to chronic health conditions. The predominance of hospitalisations for chronic conditions in this period can be primarily attributed to the large number of admissions for diabetes complications. Dehydration and gastroenteritis; and dental conditions have the highest rates of avoidable hospitalisations for the acute conditions.

Figure 8: Avoidable hospitalisations¹ by condition, North East Valley DGP and Victoria, 2001/02



¹ Admissions resulting from ACS conditions: excludes nutritional deficiencies as less than ten admissions

Table 7: Avoidable hospitalisations¹ by condition, North East Valley DGP, Victoria and Australia, 2001/02

Sub-category/ condition	North East Valley DGP		Victoria		Australia	
	No.	Rate ²	No.	Rate ²	No.	Rate ²
Vaccine-preventable	136	57.8	3,293	68.0	16,573	85.4
Influenza and pneumonia	98	41.8	2,525	52.0	13,021	67.1
Other vaccine preventable	38	16.0	768	16.0	3,552	18.3
Chronic³	3,746	1,614.6	97,133	1,982.6	352,545	1,816
Diabetes complications	1,754	756.2	44,409	906.9	141,345	728.1
Iron deficiency anaemia	253	106.7	5,196	105.9	16,451	84.7
Hypertension	59	25.0	1,362	27.7	6,354	32.7
Congestive heart failure	443	191.5	11,655	234.1	42,447	218.6
Angina	410	176.8	12,285	250.4	49,963	257.4
Chronic obstructive pulmonary disease	479	209.0	12,850	260.7	54,853	282.6
Asthma	348	149.4	9,376	196.9	41,009	211.3
Acute	2,106	891.1	50,153	1,041.7	200,913	1,035
Dehydration and gastroenteritis	500	208.7	9,761	200.0	37,766	194.5
Convulsions and epilepsy	346	146.7	7,297	152.4	31,137	160.4
Ear, nose and throat infections	262	113.8	6,653	140.5	32,075	165.2
Dental conditions	468	199.7	12,235	256.7	43,667	224.9
Perforated/bleeding ulcer	67	28.8	1,618	32.9	5,795	29.9
Ruptured appendix	39	16.4	855	17.9	3,866	19.9
Pyelonephritis	70	28.9	1,948	40.2	7,386	38.0
Pelvic inflammatory disease	60	23.6	1,693	34.8	6,547	33.7
Cellulitis	238	100.5	6,751	139.0	28,204	145.3
Gangrene	56	24.0	1,342	27.3	4,470	23.0
Total avoidable hospitalisations⁴	5,766	2,469.2	145,135	2,983.2	552,786	2,847.5

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

³ Excludes nutritional deficiencies as less than ten admissions

⁴ Sub-category and condition numbers and rates do not add to the reported total avoidable admissions: five conditions (influenza & pneumonia, other vaccine preventable, diabetes complications, ruptured appendix and gangrene) are counted in 'any diagnosis', so may be included in more than one condition group

Avoidable mortality

Avoidable and amenable mortality comprises those causes of death that are potentially avoidable at the present time, given available knowledge about social and economic policy impacts, health behaviours, and health care (the latter relating to the subset of amenable causes).

For information on the avoidable and amenable mortality conditions and ICD codes included in the analysis in this section, please refer to the *Australian and New Zealand Atlas of Avoidable Mortality*, available from www.publichealth.gov.au.

Almost three quarters (70.9%) of all deaths in North East Valley DGP at ages 0 to 74 years over the period 1997 to 2001 are considered to be avoidable, consistent with the proportion for Melbourne (71.0%) (Table 8). Deaths amenable to health care (amenable mortality, a subset of avoidable mortality) accounted for 29.4% of all deaths at ages 0 to 74 years in North East Valley DGP, compared to 28.7% in Melbourne.

Table 8: Avoidable and unavoidable mortality (0 to 74 years) by area, North East Valley DGP, Melbourne, Victoria and Australia, 1997 to 2001

Mortality category	North East Valley DGP		Melbourne		Victoria		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable	2,034	189.5	30,654	193.0	45,466	201.3	189,845	211.8
% of total	70.9	..	71.0	..	70.9	..	71.5	..
(Amenable)	(843)	(78.8)	(12,406)	(78.4)	(18,406)	(81.4)	(76,249)	(85.1)
(% of total)	(29.4)	(..)	(28.7)	(..)	(28.7)	(..)	(28.7)	(..)
Unavoidable	834	78.1	12,517	79.1	18,617	82.4	75,582	84.3
% of total	29.1	..	29.0	..	29.1	..	28.5	..
Total mortality	2,868	267.6	51,477	272.1	64,083	283.7	265,427	296.1
%	100.0	..	100.0	..	100.0	..	100.0	..

¹ Rate is the indirectly age-standardised rate per 100,000 population

Rates of avoidable mortality were higher for males than for females in each of the comparator areas. North East Valley DGP's rate of avoidable mortality for males was 241.3 deaths per 100,000 males, higher than the rate of 136.9 for females. Similarly, the rate of amenable mortality for males in the Division was higher, 83.3, compared to 74.2 for females, a rate ratio of 1.12 (Figure 9, Table 9).

Figure 9: Avoidable and amenable mortality by sex (0 to 74 years), North East Valley DGP, Melbourne, Victoria and Australia, 1997 to 2001

Note: the different scales




Table 9: Avoidable and amenable mortality (0 to 74 years) by sex, North East Valley DGP, Melbourne, Victoria and Australia, 1997 to 2001

Mortality category and sex	North East Valley DGP		Melbourne		Victoria		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
Males	1,292	241.3	19,378	244.5	29,042	257.0	123,026	272.6
Females	741	136.9	11,276	140.7	16,424	144.8	66,819	150.1
Total	2,034	189.5	30,354	193.0	45,466	201.3	189,845	211.8
Rate ratio-M:F²	..	1.76**	..	1.74**	..	1.77**	..	1.82**
Amenable								
Males	442	83.3	6,667	84.9	10,052	88.9	42,568	94.3
Females	401	74.2	5,739	71.8	8,354	73.7	33,681	75.7
Total	843	78.8	12,406	78.4	18,406	81.4	76,249	85.1
Rate ratio-M:F²	..	1.12**	..	1.18**	..	1.21**	..	1.25**

¹ Rate is the indirectly age-standardised rate per 100,000 population

² Rate ratio (M:F) is the ratio of male to female rates; rate ratios differing significantly from 1.0 are shown with * p < 0.05; ** p < 0.01

Another way of measuring premature mortality is to calculate the number of years of life lost (YLL)¹, which takes into account the years a person could have expected to live at each age of death based on the average life expectancy at that age.

The numbers of YLL for North East Valley DGP, Melbourne, Victoria and Australia over the period of analysis are shown in Table 10 by mortality category. However, given the substantial variation in the populations of these areas, a comparison of the proportion of YLL for each area is also shown.

YLL from avoidable mortality accounted for 71.7% of total YLL (0 to 74 years) for North East Valley DGP, consistent with the proportion for Melbourne. The proportion of YLL from amenable mortality for North East Valley DGP (29.1%) was slightly higher than for Melbourne (28.1%).

Table 10: Years of life lost from avoidable mortality (0 to 74 years), North East Valley DGP, Melbourne, Victoria and Australia, 1997 to 2001

Mortality category	North East Valley DGP		Melbourne		Victoria		Australia	
	No.	% of total	No.	% of total	No.	% of total	No.	% of total
Avoidable	35,602	71.7	536,388	71.6	790,054	71.5	3,327,375	71.9
(Amenable)	(14,427)	(29.1)	(210,627)	(28.1)	(310,758)	(28.1)	(1,298,430)	(28.0)
Unavoidable	14,046	28.3	212,979	28.4	315,555	28.5	1,303,289	28.1
Total	49,649	100.0	749,368	100.0	1,105,610	100.0	4,630,664	100.0

¹ Years of life lost were calculated using the remaining life expectancy method (this provides an estimate of the average time a person would have lived had he or she not died prematurely). The reference life table was the Coale and Demeny Model Life Table West level 26 female (for both males and females), with the YLL discounted to net present value at a rate of 3 per cent per year.

In each of the areas in Table 11, the majority of avoidable mortality at ages 0 to 74 years occurred in the 65 to 74 year age group (Table 11), with 1,284.4 deaths per 100,000 population in the North East Valley Division. The 45 to 64 year age group accounted for the next highest rate of avoidable death in all of the comparators, with a rate 255.6 in the North East Valley Division.

Table 11: Avoidable and amenable mortality by age, North East Valley DGP, Melbourne, Victoria and Australia, 1997 to 2001

Mortality category and age (years)	North East Valley DGP		Melbourne		Victoria		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
0-14	62	27.6	874	26.0	1,290	27.1	5,669	28.8
15-24	69	41.2	1,120	45.2	1,627	49.3	7,045	52.8
25-44	277	74.9	4,090	75.6	5,705	78.9	24,356	83.9
45-64	665	255.6	10,123	273.0	15,004	286.9	64,282	304.9
65-74	960	1,284.4	14,447	1265.1	21,840	1306.6	88,493	1,358.1
Total	2,034	189.5	30,654	193.0	45,466	201.3	189,845	211.8
Amenable								
0-24	62	16.1	836	14.6	1,189	14.9	5,083	15.4
25-44	71	19.1	963	18.0	1,382	19.1	5,946	20.5
45-64	292	111.8	4,398	118.2	6,489	123.8	27,464	130.3
65-74	418	558.3	6,209	542.7	9,348	558.6	37,756	579.4
Total	843	78.8	12,406	78.4	18,406	81.4	76,249	85.1

¹ Rate is the indirectly age-standardised rate per 100,000 population

Table 12 shows the number and age-standardised death rate by selected major condition group and selected causes included in the avoidable mortality classification.

The highest rates of avoidable mortality for the selected major condition groups in the North East Valley DGP were for cancer, with a rate of 67.9 deaths per 100,000 population, and cardiovascular diseases, 54.2 deaths per 100,000 population (Table 12, Figure 10). For the selected causes within the condition groups, the two major causes of avoidable mortality were ischaemic heart disease and lung cancer, with rates of 36.8 per 100,000 population and 21.1 per 100,000, respectively.


Table 12: Avoidable mortality (0 to 74 years) by major condition group and selected cause, North East Valley DGP, Melbourne, Victoria and Australia, 1997 to 2001

Condition group/ selected cause	North East Valley DGP		Melbourne		Victoria		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Cancer	728	67.9	10,739	67.9	15,813	69.8	62,338	69.5
Colorectal cancer	162	15.2	2,218	14.1	3,351	14.8	13,008	14.5
Lung cancer	223	21.1	3,505	22.3	5,244	23.1	21,208	23.7
Cardiovascular diseases	573	54.2	8,946	56.8	13,612	60.0	59,945	66.9
Ischaemic heart disease	389	36.8	6,377	40.6	9,809	43.3	43,712	48.8
Cerebrovascular diseases	148	13.9	2,013	12.7	2,947	12.9	12,558	14.0
Respiratory system diseases	113	10.8	1,644	10.4	2,621	11.5	11,612	13.0
Chronic obstructive pulmonary disease	96	9.2	1,451	9.2	2,339	10.2	10,395	11.6
Unintentional injuries	159	14.3	2,394	14.6	3,536	15.9	14,224	15.9
Road traffic injuries	79	7.1	1,192	7.3	1,931	8.7	8,138	9.1
Intentional injuries	140	12.5	2,074	12.6	3,020	13.6	13,891	15.5
Suicide and self inflicted injuries	132	11.8	1,877	11.4	2,752	12.3	12,393	13.8

¹ Rate is the indirectly age-standardised rate per 100,000 population

Rates in the Division were generally below or consistent with those for Melbourne and Australia for all the condition groups and selected causes (Figure 10).

Figure 10: Avoidable mortality (0 to 74 years) by major condition group and selected cause, North East Valley DGP, Melbourne and Australia, 1997 to 2001



Notes on the data

Data sources and limitations

General

References to 'Melbourne' relate to the Melbourne Statistical Division.

Data sources

Table 13 details the data sources for the material presented in this profile.

Table 13: Data sources

Section	Source
Population	
Figures 1 and 2; Table 1	Estimated Resident Population, ABS, 30 June for the periods shown
Figure 3	Estimated Resident Population, ABS, 30 June 2005; Population Projections, ABS, 30 June 2020 (unpublished) ¹
Additional socio-demographic indicators	
Figure 4	ABS SEIFA package, Census 2001
Table 2; Figure 5; Map 1	Jobless families, ABS, 2001 (unpublished)
Table 2; Figure 5; Map 2	Private health insurance, from Hansard
GP services – patient flow/ GP catchment	
Tables 3 and 4	Medicare Australia, 2003/04
Additional prevalence estimates: chronic diseases and risk factors combined	
Figure 6; Table 5	Estimated from 2001 National Health Survey (NHS), ABS (unpublished)
Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions	
Tables 6 and 7; Figures 7 and 8	National Hospital Morbidity Database at Australian Institute of Health & Welfare, 2001/02; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)
Avoidable mortality	
Tables 8, 9, 10, 11 and 12; Figures 9 and 10	ABS Deaths 1997-2001; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)

¹ The projected population at June 2020 is based on the 2002 ERP. As such, it is somewhat dated, and does not take into account more recent demographic trends: it is however the only projection series available at the SLA level for the whole of Australia.

Methods

For background information on the additional prevalence estimates presented in this profile, please refer to the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Please also refer to the November 2005 profile for information on the data converters.

Mapping

In some Divisions the maps may include a very small part of an SLA which has not been allocated any population; or has a population of less than 100 or has less than 1% of the SLAs total population; or there were less than five cases (i.e. jobless families, people with health insurance): these areas are mapped with a pattern.

Statistical geography of the North East Valley DGP

For information on the postcodes in the Division, please refer the Department of Health and Ageing website <http://www.health.gov.au/internet/wcms/publishing.nsf/Content/health-pcd-programs-divisions-divspc.htm>; also included in table format in the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Statistical Local Areas (SLAs) are defined by the Australian Bureau of Statistics to produce areas for the presentation and analysis of data. In this Division, some Local Government Areas (LGAs) have been split into SLAs. For example, the LGA of Darebin has two SLAs, Northcote (all of which is in the Division) and Preston (a small portion of which is in the Division). These SLAs and all or parts of the other SLAs listed in Table 14 comprise the Division.

Table 14: SLAs and population in North East Valley DGP, 2005 on 2001 boundaries

SLA code	SLA name	Per cent of the SLA's population in the Division *	Estimate of the SLA's 2005 population in the Division
20661	Banyule - Heidelberg	100.0	62,213
20662	Banyule - North	91.9	50,774
21892	Darebin - Preston	3.8	45,930
24214	Manningham - West	5.7	3,073
25622	Murrindindi - West	21.4	5,561
25713	Nillumbik - South	90.6	1,643
25715	Nillumbik - South West	100.0	25,613
25718	Nillumbik Balance	86.6	23,399
27074	Whittlesea - South	4.4	8,110
27351	Yarra - North	5.1	4,679

* Proportions are approximate and are known to be incorrect in some cases, due to errors in the concordance used to allocate CDs to form postal areas

Acknowledgements

Funding for these profiles was provided by the Population Health Division of the Department of Health and Ageing (DoHA).

Further developments and updates

When the re-aligned boundaries are released and DoHA have made known their geographic composition, PHIDU will examine the need to revise and re-publish these profiles (*Population health profile*, dated November 2005, and the *Population health profile: supplement*, dated March 2007).

PHIDU contact details

For general comments, data issues or enquiries re information on the web site, please contact PHIDU:

Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au