

Population health profile of the Adelaide Central and Eastern Division of General Practice: supplement

Population Profile Series: No. 89a

PHIDU

March 2007



PHIDU



Australian Government

Australian Institute of
Health and Welfare



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA



© Commonwealth of Australia 2007

This work may be reproduced and used subject to acknowledgement of the source of any material so reproduced.

National Library of Australia Cataloguing in Publication entry

Population health profile of the Adelaide Central & Eastern Division of General Practice: supplement.

Bibliography.

ISBN 9 78073089 6852 (web).

1. Public health - South Australia - Adelaide - Statistics.
2. Health status indicators - South Australia - Adelaide - Statistics.
3. Health service areas - South Australia - Adelaide.
4. Adelaide (S. Aust.) - Statistics, Medical. I. Public Health Information Development Unit (Australia). (Series : Population profile series ; no. 89a).

362.10994231

ISSN 1833-0452 Population Profile Series

Public Health Information Development Unit, The University of Adelaide
A Collaborating Unit of the Australian Institute of Health and Welfare

This profile was produced by PHIDU, the Public Health Information Development Unit at The University of Adelaide, South Australia. The work was funded under a grant from the Australian Government Department of Health and Ageing. The views expressed in this profile are solely those of the authors and should not be attributed to the Department of Health and Ageing or the Minister for Health and Ageing.

Interpretation of differences between data in this profile and similar data from other sources needs to be undertaken with care, as such differences may be due to the use of different methodology to produce the data.

Suggested citation:

PHIDU. (2007) *Population health profile of the Adelaide Central and Eastern Division of General Practice: supplement*. Population Profile Series: No. 89a. Public Health Information Development Unit (PHIDU), Adelaide.

Enquiries about or comments on this publication should be addressed to:

PHIDU, The University of Adelaide, South Australia 5005
Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au

This publication, the maps and supporting data, together with other publications on population health, are available from the PHIDU website (www.publichealth.gov.au).

Published by Public Health Information Development Unit, The University of Adelaide

Contributors: Anthea Page, Sarah Ambrose, Kristin Leahy and John Glover

Population health profile of the Adelaide Central and Eastern Division of General Practice: supplement

This profile is a supplement to the *Population health profile of the Adelaide Central and Eastern Division of General Practice*, dated November 2005, available from www.publichealth.gov.au. This supplement includes an update of the population of the Adelaide Central & Eastern Division of General Practice, as well as additional indicators and aspects of the Division's socioeconomic status, use of GP services and health. The contents are:


- Population [updated to June 2005]
- Additional socio-demographic indicators
- Unreferred attendances – patient flow/ GP catchment
- Additional prevalence estimates: chronic diseases and risk factors combined
- Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions
- Avoidable mortality

For further information on the way Division totals in this report have been estimated, please refer to the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Population

The Adelaide Central and Eastern Division had an Estimated Resident Population of 182,072 at 30 June 2005.

Figure 1: Annual population change, Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 1991 to 1996, 1996 to 2001 and 2001 to 2005




Over the five years from 1991 to 1996, the Division's population increased by 0.3% on average each year, below the rate for Adelaide and South Australia (both 0.4%). From 1996 to 2001, the annual percentage increase in the Division was 0.4%, lower than the increases for Adelaide and South Australia (both 0.6%). The Division's growth rate decreased to 0.2% from 2001 to 2005, again lower than the annual increases for Adelaide and South Australia (both 0.4%). Population change at the national level was much higher in all periods.

Table 1: Population by age, Adelaide Central & Eastern DGP and Australia, 2005

Age group (years)	Adelaide Central & Eastern DGP		Australia	
	No.	%	No.	%
0-14	26,130	14.4	3,978,221	19.6
15-24	27,428	15.1	2,819,834	13.9
25-44	50,400	27.7	5,878,107	28.9
45-64	46,354	25.5	4,984,446	24.5
65-74	14,358	7.9	1,398,831	6.9
75-84	12,170	6.7	954,143	4.7
85+	5,232	2.9	315,027	1.5
Total	182,072	100.0	20,328,609	100.0

As shown in the accompanying table and the age-sex pyramid (Figure 2), the Adelaide Central and Eastern DGP had notably fewer children than Australia as a whole, with 14.4% at ages 0 to 14 years (compared to 19.6% for Australia) (Table 1). Conversely, the proportions of the Division's population aged 45 years and over were higher, in some cases substantially so, than for Australia.


Figure 2: Population in Adelaide Central & Eastern DGP and Australia, by age and sex, 2005



The most notable differences in the age distribution of the Division's population (when compared to Australia overall) are:

- at younger ages – substantially fewer children aged 0 to 14 years;
- from 20 to 24 years – substantially more males and females;
- from 30 to 44 years – relatively fewer females; and
- at older ages – relatively more males and females aged 55 years and over (most notably, females at ages 75 years and over).

Figure 3: Population projections for Adelaide Central & Eastern DGP, by age and sex, 2005 and 2020




The population projections for the Division show a number of changes in age distribution, with the 2020 population projected to have:

- at younger ages – relatively fewer children, young people and young adults, aged 0 to 24 years;
- from 30 to 49 years – relatively fewer males and females; and
- from age 60 onwards – relatively more males and females (excluding 80 to 84 year old females).

Additional socio-demographic indicators

Please refer to the earlier *Population health profile of the Adelaide Central and Eastern Division of General Practice*, dated November 2005, available from www.publichealth.gov.au, for other socio-demographic indicators.

Figure 4: Index of Relative Socio-Economic Disadvantage, Adelaide Central & Eastern DGP, 2001



One of four socioeconomic indexes for areas produced at the 2001 ABS Census is the Index of Relative Socio-Economic Disadvantage.

The Adelaide Central and Eastern DGP has an index score of 1071, above the score for Australia of 1000: this score varies across the Division, from 980 in the most disadvantaged areas to 1140 in the least disadvantaged areas.


Note: each 'quintile' comprises approximately 20% of the population of the Division.

A new indicator, produced for the first time at the 2001 ABS Census, shows the number of jobless families with children under 15 years of age. There were markedly fewer jobless families in the Adelaide Central and Eastern DGP (11.6%), compared to Adelaide as a whole (19.2%) (Figure 5, Table 2).

With the introduction of the 30% rebate for private health insurance premiums, there was a once-off registration process, providing information of the postcode and residence of those who had such insurance (these data are not available at this area level for later dates). In 2001, the Division had a markedly higher proportion of people with private health insurance (63.5%), compared to Adelaide (54.2%) (Figure 5, Table 2).

Figure 5: Socio-demographic indicators, Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 2001

Jobless families with children under 15 years old



Private health insurance, 30 June





Table 2: Socio-demographic indicators, Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 2001

Indicator	Adelaide Central & Eastern DGP		Adelaide		South Australia		Australia	
	No.	%	No.	%	No.	%	No.	%
Jobless families with children under 15 years old	1,708	11.6	21,478	19.2	29,203	18.7	357,563	17.4
Private health insurance (30 June)	113,069	63.5	581,532	54.2	754,598	51.3	8,671,106	46.0

Details of the distribution of jobless families (Map 1) and of the population covered by private health insurance (Map 2) are shown by Statistical Local Area (SLA) in Maps 1 and 2, respectively.


Map 1: Jobless families with children under 15 years of age by SLA, Adelaide Central & Eastern DGP, 2001



[#] data were not mapped: see 'Mapping' note under Methods

For map labels: see next page


Map 2: People covered by private health insurance by SLA, Adelaide Central & Eastern DGP, 30 June 2001



Map legend for SLAs not distinguishable

- Tea Tree Gully - South

Per cent



data were not mapped: see 'Mapping' note under Methods

Alphabetical key to Statistical Local Areas, Adelaide Central & Eastern DGP, 2001

Adelaide	4	Norwood Payneham St Peters - East	10
Adelaide Hills - Ranges	5	Norwood Payneham St Peters -West	11
Burnside - North-East	6	Tea Tree Gully - South	1
Burnside - South-West	7	Unley - East	12
Campbelltown - East	8	Unley - West	13
Campbelltown - West	9	Walkerville	14
Marion - North	15	West Torrens - East	2
Mitcham - North-East	16	West Torrens - West	3

GP services to residents of the Adelaide Central and Eastern DGP

The following tables include information, purchased from Medicare Australia, of the movement of patients and GPs between Divisions. Note that the data only include unreferral attendances recorded under Medicare: unreferral attendances not included are those for which the cost is met by the Department of Veterans' Affairs or a compensation scheme; or are provided by salaried medical officers in hospitals, community health services or Aboriginal Medical Services, and which are not billed to Medicare. At any attendance, one or more services may have been provided.

Over two thirds (70.7%) of unreferral attendances to residents of Adelaide Central and Eastern DGP were provided in the Division (ie. by a GP with a provider number in the Division): this represented 623,807 GP unreferral attendances (Table 3). A further 10.2% of unreferral attendances to residents of the Division were provided by GPs in Adelaide North East DGP, with approximately 7.0% by GPs in each of Southern DGP and Adelaide Western DGP.

Table 3: Patient flow – People living¹ in Adelaide Central and Eastern DGP by Division where attendance occurred², 2003/04

Division		Unreferral attendances	
Number	Name	No.	% ³
504	Adelaide Central and Eastern DGP	623,807	70.7
503	Adelaide North East DGP	89,991	10.2
505	Southern DGP	67,950	7.7
501	Adelaide Western DGP	61,919	7.0
514	Adelaide Hills DGP	10,518	1.2
Other	..	28,541	3.2
Total	..	882,726	100.0

¹ Based on address in Medicare records

² Division of GP based on provider number

³ Proportion of all unreferral attendances of patients with an address in Division 504 by Division in which attendance occurred

Just over half (55.1%) of unreferral attendances provided by GPs with a provider number in Adelaide Central and Eastern DGP were to people living in the Division (ie. their Medicare address was in the Division) (Table 4). A further 12.7% of unreferral attendances were provided to residents of Southern DGP, with 12.3% from North East DGP and 10.9% provided to people from Adelaide Western DGP. The relatively low proportion of patients represented by residents of the Division (55.1%) reflects the relatively large number of people coming into the Division during the day to work (in particular in the city centre), some of whom use GPs in the Division.

Table 4: GP catchment – Unreferral attendances provided by GPs¹ in Adelaide Central and Eastern DGP by Division of patient address², 2003/04

Division		Unreferral attendances	
Number	Name	No.	% ³
504	Adelaide Central and Eastern DGP	623,807	55.1
505	Southern DGP	143,838	12.7
503	Adelaide North East DGP	139,351	12.3
501	Adelaide Western DGP	123,915	10.9
502	Adelaide Northern DGP	48,818	4.3
514	Adelaide Hills DGP	17,742	1.6
Other	..	35,527	2.3
Total	..	1,132,998	100.0

¹ Division of GP based on provider number

² Based on address in Medicare records

³ Proportion of all unreferral attendances to GPs with a provider number in Division 504 by Division of patient address

Additional prevalence estimates: chronic diseases and risk factors combined

Please refer to the earlier *Population health profile of the Adelaide Central and Eastern Division of General Practice*, dated November 2005, available from www.publichealth.gov.au, for the separate prevalence estimates of chronic disease; measures of self-reported health and risk factors. The process by which the estimates have been made, and details of their limitations, are also described in the 'Notes on the data' section of this earlier profile.

In this section two estimates, which combine the prevalence of selected chronic diseases with a risk factor, are shown for the Division. The measures are of people who *had asthma and were smokers*, and people who *had type 2 diabetes and were overweight or obese*: note that the estimates have been predicted from self-reported data, and are not based on clinical records or physical measures.

It is estimated that there were markedly fewer people in Adelaide Central and Eastern DGP who had asthma and were smokers, compared to Adelaide or Australia as a whole (Figure 6, Table 5): that is, the prevalence rates per 1,000 population were lower. The rate of people in Adelaide Central and Eastern DGP who had type 2 diabetes and were overweight/ obese was slightly lower than in Adelaide or Australia.

Figure 6: Estimates of selected chronic diseases and risk factors, Adelaide Central & Eastern DGP, Adelaide and Australia, 2001




Table 5: Estimates of selected chronic diseases and risk factors, Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 2001

Variable	Adelaide Central & Eastern DGP		Adelaide		South Australia		Australia	
	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ¹
Had asthma & smoked ³	3,114	16.4	23,430	21.3	32,487	22.3	397,734	20.8
Had type 2 diabetes & were overweight/ obese ⁴	2,939	14.5	17,762	15.7	23,187	14.9	283,176	15.2

¹ No. is a weighted estimate of the number of people in Adelaide Central & Eastern DGP reporting these chronic conditions/ with these risk factors and is derived from synthetic predictions from the 2001 NHS

² Rate is the indirectly age-standardised rate per 1,000 population

³ Population aged 18 years and over

⁴ Population aged 15 years and over

Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions

The rationale underlying the concept of avoidable hospitalisations is that timely and effective care of certain conditions, delivered in a primary care setting, can reduce the risk of hospitalisation. Admissions to hospital for these ambulatory care sensitive (ACS) conditions can be avoided in three ways. Firstly, for conditions that are usually preventable through immunisation or nutritional intervention, disease can be prevented almost entirely. Secondly, diseases or conditions that can lead to rapid onset problems, such as dehydration and gastroenteritis, can be treated. Thirdly, chronic conditions, such as congestive heart failure, can be managed to prevent or reduce the severity of acute flare-ups to avoid hospitalisation.

This measure does not include other aspects of avoidable morbidity, namely potentially preventable hospitalisations (hospitalisations resulting from diseases preventable through population based health promotion strategies, e.g. alcohol-related conditions; and most cases of lung cancer) and hospitalisations avoidable through injury prevention (e.g. road traffic accidents).

For information on the ambulatory care sensitive conditions and ICD codes included in the analysis in this section, please refer to the *Atlas of Avoidable Hospitalisations in Australia: ambulatory care-sensitive conditions*, available from www.publichealth.gov.au.

In 2001 to 2002, the 4,612 admissions from ambulatory care sensitive (ACS) conditions accounted for 7.4% of all admissions in the Adelaide Central and Eastern DGP (Table 6, Figure 7), notably below the levels in South Australia (8.5) and Australia (8.7%).


Table 6: Avoidable¹ and unavoidable hospitalisations, Adelaide Central & Eastern DGP, South Australia, and Australia, 2001/02

Category	Adelaide Central & Eastern DGP			South Australia			Australia		
	No.	Rate ²	%	No.	Rate ²	%	No.	Rate ²	%
Avoidable ¹	4,612	2,169.2	7.4	47,247	2,915.7	8.5	552,786	2,847.5	8.7
Unavoidable	57,608	28,397.2	92.6	507,053	32,039.4	91.5	5,818,199	29,970.7	91.3
Total	62,220	30,543.3	100.0	554,300	34,952.2	100.0	6,370,985	32,818.2	100.0

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

Figure 7: Avoidable hospitalisations¹, Adelaide Central & Eastern DGP, South Australia and Australia, 2001/02




The rate of avoidable hospitalisations in Adelaide Central and Eastern DGP is markedly lower, a rate of 2,169.2 admissions per 100,000 population, compared to both South Australia (a rate of 2,915.7) and Australia (2,847.5).

¹ Admissions resulting from ACS conditions

Diabetes complications, dental conditions and asthma were the three conditions with the highest rates of avoidable hospitalisations in the Adelaide Central and Eastern DGP (Figure 8, Table 7).

Table 7 shows the number, rate and proportion of avoidable hospitalisations, for the individual ACS conditions, as well as the vaccine-preventable; acute; and chronic sub-categories. Two-thirds of avoidable hospitalisations are attributable to chronic health conditions. The predominance of hospitalisations for chronic conditions in this period can be primarily attributed to the large number of admissions for diabetes complications. Dental conditions; and ear, nose and throat infections have the highest rates of avoidable hospitalisations for the acute conditions.

Figure 8: Avoidable hospitalisations¹ by condition, Adelaide Central & Eastern DGP and South Australia, 2001/02



¹ Admissions resulting from ACS conditions: excludes nutritional deficiencies as less than ten admissions

Table 7: Avoidable hospitalisations¹ by condition, Adelaide Central & Eastern DGP, South Australia and Australia, 2001/02

Sub-category/ condition	Adelaide Central & Eastern DGP		South Australia		Australia	
	No.	Rate ²	No.	Rate ²	No.	Rate ²
Vaccine-preventable	179	89.5	1,466	92.9	16,573	85.4
Influenza and pneumonia	128	61.1	1,075	67.0	13,021	67.1
Other vaccine preventable	51	28.4	391	25.9	3,552	18.3
Chronic³	2,862	1,275.4	30,607	1,837.6	352,545	1,816
Diabetes complications	984	435.3	11,640	692.9	141,345	728.1
Iron deficiency anaemia	161	69.7	1,271	76.1	16,451	84.7
Hypertension	35	14.7	532	31.6	6,354	32.7
Congestive heart failure	480	172.4	3,900	219.1	42,447	218.6
Angina	257	107.8	3,778	221.6	49,963	257.4
Chronic obstructive pulmonary disease	522	217.6	4,710	272.9	54,853	282.6
Asthma	423	257.9	4,776	323.4	41,009	211.3
Acute	1,668	921.0	16,405	1,077.6	200,913	1,035
Dehydration and gastroenteritis	294	137.6	3,111	194.8	37,766	194.5
Convulsions and epilepsy	192	109.7	2,153	143.6	31,137	160.4
Ear, nose and throat infections	286	186.6	3,046	210.9	32,075	165.2
Dental conditions	442	269.6	3,831	259.2	43,667	224.9
Perforated/bleeding ulcer	66	26.8	555	32.5	5,795	29.9
Ruptured appendix	19	10.6	255	17.0	3,866	19.9
Pyelonephritis	93	47.9	681	44.7	7,386	38.0
Pelvic inflammatory disease	36	20.0	497	33.7	6,547	33.7
Cellulitis	214	101.3	1,987	124.1	28,204	145.3
Gangrene	26	10.9	289	17.1	4,470	23.0
Total avoidable hospitalisations⁴	4,612	2,169.2	47,247	2,915.7	552,786	2,847.5

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

³ Excludes nutritional deficiencies as less than ten admissions

⁴ Sub-category and condition numbers and rates do not add to the reported total avoidable admissions: five conditions (influenza & pneumonia, other vaccine preventable, diabetes complications, ruptured appendix and gangrene) are counted in 'any diagnosis', so may be included in more than one condition group

Avoidable mortality

Avoidable and amenable mortality comprises those causes of death that are potentially avoidable at the present time, given available knowledge about social and economic policy impacts, health behaviours, and health care (the latter relating to the subset of amenable causes).

For information on the avoidable and amenable mortality conditions and ICD codes included in the analysis in this section, please refer to the *Australian and New Zealand Atlas of Avoidable Mortality*, available from www.publichealth.gov.au.

Over two thirds (69.5%) of all deaths in Adelaide Central and Eastern DGP at ages 0 to 74 years over the period 1997 to 2001 are considered to be avoidable, slightly lower than the proportion for Adelaide (71.0%) (Table 8). However, the rate in the Division is notably lower than that in Adelaide, a differential of 0.89.

Deaths amenable to health care (amenable mortality, a subset of avoidable mortality) accounted for 29.0% of all deaths at ages 0 to 74 years in Adelaide Central and Eastern DGP, compared to 29.2% in Adelaide.

Table 8: Avoidable and unavoidable mortality (0 to 74 years) by area, Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 1997 to 2001

Mortality category	Adelaide Central & Eastern DGP		Adelaide		South Australia		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable	1,685	181.5	11,086	202.8	15,938	210.4	189,845	211.8
% of total	69.5	..	71.0	..	71.4	..	71.5	..
(Amenable)	(703)	(75.6)	(4,563)	(82.9)	(6,556)	(85.9)	(76,249)	(85.1)
(% of total)	(29.0)	(..)	(29.2)	(..)	(29.4)	(..)	(28.7)	(..)
Unavoidable	742	79.9	4,532	82.6	6,369	83.7	75,582	84.3
% of total	30.6	..	29.0	..	28.6	..	28.5	..
Total mortality	2,426	261.4	15,619	285.4	22,307	294.1	265,427	296.1
%	100.0	..	100.0	..	100.0	..	100.0	..

¹ Rate is the indirectly age-standardised rate per 100,000 population

Rates of avoidable mortality were higher for males than for females in each of the comparator areas. Adelaide Central and Eastern DGP's rate of avoidable mortality for males was 234.5 deaths per 100,000 males, higher than the rate of 127.8 for females. Similarly, the rate of amenable mortality for males in the Division was higher, 84.0, compared to 67.0 for females, a rate ratio of 1.25 (Figure 9, Table 9).

Figure 9: Avoidable and amenable mortality by sex (0 to 74 years), Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 1997 to 2001

Note: the different scales




Table 9: Avoidable and amenable mortality (0 to 74 years) by sex, Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 1997 to 2001

Mortality category and sex	Adelaide Central & Eastern DGP		Adelaide		South Australia		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
Males	1,070	234.5	7,067	261.5	10,326	272.8	123,026	272.6
Females	615	127.8	4,019	143.3	5,612	147.2	66,819	150.1
Total	1,685	181.5	11,086	202.8	15,938	210.4	189,845	211.8
Rate ratio–M:F²	..	1.83**	..	1.82**	..	1.85**	..	1.82**
Amenable								
Males	383	84.0	2,503	92.1	3,671	96.0	42,568	94.3
Females	320	67.0	2,060	73.6	2,884	75.7	33,681	75.7
Total	703	75.6	4,563	82.9	6,556	85.9	76,249	85.1
Rate ratio–M:F²	..	1.25*	..	1.25**	..	1.27**	..	1.25**

¹ Rate is the indirectly age-standardised rate per 100,000 population

² Rate ratio (M:F) is the ratio of male to female rates; rate ratios differing significantly from 1.0 are shown with * p < 0.05; ** p < 0.01

Another way of measuring premature mortality is to calculate the number of years of life lost (YLL)¹, which takes into account the years a person could have expected to live at each age of death based on the average life expectancy at that age.

The numbers of YLL for Adelaide Central and Eastern DGP, Adelaide, South Australia and Australia over the period of analysis are shown in Table 10 by mortality category. However, given the substantial variation in the populations of these areas, a comparison of the proportion of YLL for each area is also shown.

YLL from avoidable mortality accounted for 70.1% of total YLL (0 to 74 years) for Adelaide Central and Eastern DGP, lower than the 71.3% for Adelaide. The proportion of YLL from amenable mortality of 28.1% for Adelaide Central and Eastern DGP was marginally lower than the 28.5% for Adelaide.

Table 10: Years of life lost from avoidable mortality (0 to 74 years), Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 1997 to 2001

Mortality category	Adelaide Central & Eastern DGP		Adelaide		South Australia		Australia	
	No.	% of total	No.	% of total	No.	% of total	No.	% of total
Avoidable	28,807	70.1	189,430	71.3	273,135	71.8	3,327,375	71.9
(Amenable)	(11,544)	(28.1)	(75,612)	(28.5)	(108,777)	(28.6)	(1,298,430)	(28.0)
Unavoidable	12,296	29.9	76,164	28.7	107,223	28.2	1,303,289	28.1
Total	41,103	100.0	265,594	100.0	380,358	100.0	4,630,664	100.0

¹ Years of life lost were calculated using the remaining life expectancy method (this provides an estimate of the average time a person would have lived had he or she not died prematurely). The reference life table was the Coale and Demeny Model Life Table West level 26 female (for both males and females), with the YLL discounted to net present value at a rate of 3 per cent per year.

In each of the areas in Table 11, the majority of avoidable mortality at ages 0 to 74 years occurred in the 65 to 74 year age group (Table 11), with 1,170.2 deaths per 100,000 population in the Adelaide Central and Eastern Division. The 45 to 64 year age group accounted for the next highest rate of avoidable death in all of the comparators, with a rate of 251.4 in the Adelaide Central and Eastern Division.

Table 11: Avoidable and amenable mortality by age, Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 1997 to 2001

Mortality category and age (years)	Adelaide Central & Eastern DGP		Adelaide		South Australia		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
0-14	30	23.0	234	23.1	352	24.2	5,669	28.8
15-24	59	42.7	364	47.7	523	52.4	7,045	52.8
25-44	213	80.2	1,383	85.0	1,979	88.8	24,356	83.9
45-64	525	251.4	3,490	283.6	5,130	297.8	64,282	304.9
65-74	858	1,170.2	5,616	1322.6	7,954	1354.8	88,493	1,358.1
Total	1,685	181.5	11,086	202.8	15,938	210.4	189,845	211.8
Amenable								
0-24	27	11.3	223	12.9	324	13.3	5,083	15.4
25-44	56	21.6	361	22.1	507	22.6	5,946	20.5
45-64	237	112.6	1,538	124.3	2,248	130.1	27,464	130.3
65-74	383	519.2	2,441	572.8	3,477	591.6	37,756	579.4
Total	703	75.6	4,563	82.9	6,556	85.9	76,249	85.1

¹ Rate is the indirectly age-standardised rate per 100,000 population

Table 12 shows the number and age-standardised death rate by selected major condition group and selected causes included in the avoidable mortality classification.

The highest rates of avoidable mortality for the selected major condition groups in the Adelaide Central and Eastern DGP were for cancer and cardiovascular diseases, both with a rate of 57.7 deaths per 100,000 population (Table 12, Figure 10). For the selected causes within the condition groups, the two major causes of avoidable mortality were ischaemic heart disease and lung cancer, with rates of 41.9 per 100,000 population and 17.2 per 100,000, respectively.


Table 12: Avoidable mortality (0 to 74 years) by major condition group and selected cause, Adelaide Central & Eastern DGP, Adelaide, South Australia and Australia, 1997 to 2001

Condition group/ selected cause	Adelaide Central & Eastern DGP		Adelaide		South Australia		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Cancer	545	57.7	3,720	67.2	5,209	67.8	62,338	69.5
Colorectal cancer	125	13.1	796	14.4	1,142	14.8	13,008	14.5
Lung cancer	164	17.2	1,251	22.5	1,728	22.3	21,208	23.7
Cardiovascular diseases	553	57.7	3,655	65.3	5,324	68.5	59,945	66.9
Ischaemic heart disease	399	41.9	2,658	47.7	3,918	50.5	43,712	48.8
Cerebrovascular diseases	129	13.3	770	13.6	1,086	13.9	12,558	14.0
Respiratory system diseases	87	8.9	627	11.1	897	11.4	11,612	13.0
Chronic obstructive pulmonary disease	74	7.5	544	9.5	783	9.9	10,395	11.6
Unintentional injuries	109	12.7	673	13.1	1,085	15.5	14,224	15.9
Road traffic injuries	54	6.3	380	7.4	687	9.9	8,138	9.1
Intentional injuries	129	14.9	809	15.8	1,138	16.3	13,891	15.5
Suicide and self inflicted injuries	120	13.9	725	14.1	1,018	14.5	12,393	13.8

¹ Rate is the indirectly age-standardised rate per 100,000 population

For all of the condition groups and selected causes, rates in the Division were below, or consistent with, those for Adelaide and Australia (Figure 10).

Figure 10: Avoidable mortality (0 to 74 years) by major condition group and selected cause, Adelaide Central & Eastern DGP, Adelaide and Australia, 1997 to 2001



Notes on the data

Data sources and limitations

General

References to 'Adelaide' relate to the Adelaide Statistical Division.

Data sources

Table 13 details the data sources for the material presented in this profile.

Table 13: Data sources

Section	Source
Population	
Figures 1 and 2; Table 1	Estimated Resident Population, ABS, 30 June for the periods shown
Figure 3	Estimated Resident Population, ABS, 30 June 2005; Population Projections, ABS, 30 June 2020 (unpublished) ¹
Additional socio-demographic indicators	
Figure 4	ABS SEIFA package, Census 2001
Table 2; Figure 5; Map 1	Jobless families, ABS, 2001 (unpublished)
Table 2; Figure 5; Map 2	Private health insurance, from Hansard
GP services – patient flow/ GP catchment	
Tables 3 and 4	Medicare Australia, 2003/04
Additional prevalence estimates: chronic diseases and risk factors combined	
Figure 6; Table 5	Estimated from 2001 National Health Survey (NHS), ABS (unpublished)
Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions	
Tables 6 and 7; Figures 7 and 8	National Hospital Morbidity Database at Australian Institute of Health & Welfare, 2001/02; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)
Avoidable mortality	
Tables 8, 9, 10, 11 and 12; Figures 9 and 10	ABS Deaths 1997-2001; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)

¹ The projected population at June 2020 is based on the 2002 ERP. As such, it is somewhat dated, and does not take into account more recent demographic trends: it is however the only projection series available at the SLA level for the whole of Australia.

Methods

For background information on the additional prevalence estimates presented in this profile, please refer to the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Please also refer to the November 2005 profile for information on the data converters.

Mapping

In some Divisions the maps may include a very small part of an SLA which has not been allocated any population; or has a population of less than 100 or has less than 1% of the SLAs total population; or there were less than five cases (ie. jobless families, people with health insurance): these areas are mapped with a pattern.

Statistical geography of the Adelaide Central & Eastern DGP

For information on the postcodes in the Division, please refer the Department of Health and Ageing website <http://www.health.gov.au/internet/wcms/publishing.nsf/Content/health-pcd-programs-divisions-divspc.htm>; also included in table format in the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Statistical Local Areas (SLAs) are defined by the Australian Bureau of Statistics to produce areas for the presentation and analysis of data. In this Division, most Local Government Areas (LGAs) have been split into SLAs. For example, Unley is comprised of two SLAs - East and West. All of these SLAs, and all or parts of the other SLAs listed in Table 14 comprise the Division.

Table 14: SLAs and population in Adelaide Central & Eastern DGP, 2005 on 2001 boundaries

SLA code	SLA name	Per cent of the SLA's population in the Division*	Estimate of the SLA's 2005 population in the Division
40070	Adelaide	100.0	14,725
40124	Adelaide Hills - Ranges	27.5	2,840
40701	Burnside - North-East	100.0	21,648
40704	Burnside - South-West	100.0	21,292
40911	Campbelltown - East	83.0	23,037
40914	Campbelltown - West	88.8	17,339
44064	Marion - North	12.9	3,279
44344	Mitcham - North-East	6.7	1,050
45291	Norwood Payneham St Peters - East	58.1	9,431
45294	Norwood Payneham St Peters - West	100.0	17,787
46510	Prospect	2.4	469
47708	Tea Tree Gully South	5.5	1,805
47981	Unley - East	100.0	19,447
47984	Unley - West	100.0	16,884
48260	Walkerville	30.0	2,123
48411	West Torrens - East	21.9	5,218
48414	West Torrens - West	12.9	3,697

* Proportions are approximate and are known to be incorrect in some cases, due to errors in the concordance used to allocate CDs to form postal areas

Acknowledgements

Funding for these profiles was provided by the Population Health Division of the Department of Health and Ageing (DoHA).

Further developments and updates

When the re-aligned boundaries are released and DoHA have made known their geographic composition, PHIDU will examine the need to revise and re-publish these profiles (*Population health profile*, dated November 2005, and the *Population health profile: supplement*, dated March 2007).

PHIDU contact details

For general comments, data issues or enquiries re information on the web site, please contact PHIDU:

Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au