

References

- Ansari, Z. 2007. The concept and usefulness of ambulatory care sensitive conditions as indicators of quality and access to primary health care. *Australian Journal of Primary Health*. 13(3): 91-110.
- Ansari, Z. Haider, S.I., Ansari, H., de Goyer, T., and Sindall, C. 2012. Patient characteristics associated with hospitalisations for ambulatory care sensitive conditions in Victoria, Australia. *BMC Health Services Research* 12:475.
- Australian Government Department of Health. 2019. Keeping Australians Out of Hospital. <https://beta.health.gov.au/initiatives-and-programs/keeping-australians-out-of-hospital> (Accessed 21/06/2019). Australian Government.
- Australian Institute of Health and Welfare (AIHW). 2018a. National Healthcare Agreement :PI 18- Selected Potentially Preventable Hospitalisations, 2018 Australian Institute of Health and Welfare : <https://meteor.aihw.gov.au/content/index.phtml/itemId/658499> (accessed 21/06/2019).
- Australian Institute of Health and Welfare (AIHW). 2018b. Admitted patient care 2016-17: Australian hospital statistics. Health services series no.84.Cat. no. HSE 201. Canberra: AIHW
- Australian Institute of Health and Welfare (AIHW). 2018c. A potentially preventable hospitalisation indicator for general practice: consultation paper. Cat. no. HSE 214. Canberra: AIHW.
- Australian Institute of Health and Welfare (AIHW). 2018d. Australia's health 2018. Australia's health series no. 16. AUS 221. Canberra: AIHW.
- Australian Institute of Health and Welfare (AIHW). 2018e. Potentially Preventable hospitalisation in Australia by small geographic areas. Canberra: AIHW. <https://www.aihw.gov.au/reports/primary-health-care/potentially-preventable-hospitalisations/contents/overview> (accessed 21/06/2019).
- Banham, D., Woolacott, T., Gray, J., Humphrys, B., Mihnev, A., McDermott, R. 2010. Recognising potential for preventing hospitalisations. *Australian Health Review*. 34: 116-122.
- Banham, D., Chen, T., Karnon, J., Brown, A. and Lynch, J. 2017. Sociodemographic variations in the amount, duration and cost of potentially preventable hospitalisation for chronic conditions among Aboriginal and non-Aboriginal Australians: a period prevalence study of linked public hospital data. *BMJ Open* 7:e017331.
- Basu, A. and Brinson, D. 2008. The effectiveness of interventions for reducing ambulatory sensitive hospitalisations: a systematic review. *HSAC Report 2008*; 1(6).
- Bird, S., Noronha, M., and Sinnott, H. 2010. An integrated care facilitation model improves quality of life and reduces use of hospital resources by patients with chronic obstructive pulmonary disease and chronic heart failure. *Australian Journal of Primary Care*. 66:326-333.
- Busby, J., Purdy, S., and Hollingworth, W. 2015. A systematic review of the magnitude and cause of geographic variation in unplanned hospital admission rates and length of stay for ambulatory care sensitive conditions. *BMC Health Services Research* 15:324.
- Busby, J., Purdy, S., and Hollingworth, W. 2017a. Using geographic variation in unplanned ambulatory care sensitive condition admission rates to identify commissioning priorities: an analysis of routine data from England. *Journal of Health Service Research and Policy*. 22(1) 20-27.

Busby, J. Purdy, S and Hollingworth, W. 2017b. How do population, general practice and hospital factors influence ambulatory care sensitive admissions: a cross sectional study. *BMC Family practice* 18:67.

Dantas, I., Santana, R., Sarmiento, J., and Aguiar, P. 2016. The impact of multiple chronic diseases on hospitalisations for ambulatory care sensitive conditions. *BMC Health Services Research* 16:348.

Department of Health Western Australia (DOHWA). 2017. Lessons of Location: Potentially preventable hospitalisation hotspots in Western Australia 2017. Government of Western Australia Department of Health. WA Primary Health Alliance. https://ww2.health.wa.gov.au/~/_media/Files/Corporate/Reports%20and%20publications/Lessons-of-Location/Lessons-of-Location-2017.pdf (accessed 26/06/2019).

Duckett, S., and Griffiths, K., 2016. Perils of place: identifying hotspots of health inequalities. Grattan Institute.

Einarsdottir, K., Preen, D.B. Emery, J.D., Kelman, C., and Holman, C.D.J. 2010. Regular primary care lowers hospitalisation risk and mortality in seniors with chronic respiratory diseases. *J Gen Intern Med* 25(8):766-773.

Erny-Albrecht K, Oliver-Baxter J, and Bywood P. 2016. Primary health care-based programmes targeting potentially avoidable hospitalisations in vulnerable groups with chronic disease. PHCRIS Policy Issue Review. Adelaide: Primary Health Care Research & Information Service.

Falster, M.O., Jorm, L.R., Douglas, K.A, Blyth, F.M., Elliott, R.F., and Leyland, A.H. 2015. Sociodemographic and health characteristics, rather than primary care supply, are major drivers of geographic variation in preventable hospitalisation in Australia. *Medical Care* 53(5).

Falster, M and Jorm, L. 2017. A guide to the potentially preventable hospitalisations indicator in Australia. Centre for Big Data Research in Health, University of New South Wales in consultation with Australian Commission on Safety and Quality in Health Care and Australian Institute of Health and Welfare: Sydney; 2017.

Falster, M.O., Leyland, A.H., and Jorm, L.R. 2019. Do hospitals influence geographic variation in admission for preventable hospitalisation? A data linkage study in New South Wales, Australia. *BMJ Open*;9:e027639.

Fleetcroft, R., Hardcastle, A., Steel, N., Price, G.M., Purdy, S., Lipp, A., Myint, P.K and Howe, A. 2018. Does practice analysis agree with the ambulatory care sensitive conditions' list of avoidable unplanned admissions? : a cross sectional study in the East of England. *BMJ Open* 8:e020756.

Freund, T., Campbell, S.M., Geissler, S., Kunz, C.U., Mabler, C., Peters-Klimm, F. and Szecsenyi, J. 2013. Strategies for reducing potentially avoidable hospitalisations for ambulatory care-sensitive conditions. *Annals of Family Medicine* 11(4).

Gavidia, T., Varhol, R., Xiao, A., Koh, C., Mai, Q., Liu, Y., Turdukulov, U., Parsons, S., Fievez, P., Veenendaal, B., and Somerford, P. 2019. Geographic variation in primary health care service utilisation and potentially preventable hospitalisations in Western Australia. Department of Health WA. <https://ww2.health.wa.gov.au/Reports-and-publications/Geographic-variation-in-primary-healthcare-service-utilisation> (Accessed 26/06/2019).

Gibson, O.R, Segal, L., and McDermott, R.A. 2013. A systematic review of evidence on the association between hospitalisation for chronic disease related ambulatory care sensitive conditions and primary health care resourcing. *BMC Health Services Research* 13:336.

Ha, N.T., Harris, M., Preen, D., Robinson, S. and Moorin, R. 2018. Identifying patterns of general practitioner service utilisation and their relationship with potentially preventable hospitalisations in people with diabetes: The utility of a cluster analysis approach. *Diabetes Research and Clinical Practice* 138: 201-210.

Harriss, L.R., Thompson, F., Lawson, K., O Loughlin, M.O., and McDermott, R. 2018. Preventable hospitalisation in regional Queensland: potential for primary health? *Australian Health Review*. Early view (<https://doi.org/10.1071/AH18033>), 1-11. <https://doi.org/10.1071/AH18033>.

Harrold, T.C., Randall, D.A., Falster, M.O., Lujic, S., and Jorm, L.R. 2014. The contribution of geography in preventable hospitalisations between Indigenous and non-indigenous Australians. *PLOS One* 9(5).

Health Performance Council (HPC). 2017. Hotspots of potentially preventable hospitalisations in South Australia's public hospitals. Pinpointing potential health inequalities by analysis of public hospital activity data. Government of South Australia. December 2017. <http://www.hpcs.com.au/statistics/potentially-preventable-hospitalisations> (accessed 26/06/2019).

Hollingworth, S.A., Donald, M., Zhang, J., Vaikuntam, B.P., Russell, A. and Jackson, C. 2017. Impact of a general practitioner-led integrated model of care on the cost of potentially preventable diabetes-related hospitalisations. *Primary Care Diabetes II*: 344-347.

Jaques, K., Bernstein, M., Haigh, F., and Zapart, S. 2018. Reducing the Risk of Potentially Preventable Hospitalisations: A Literature Review of Community-based Approaches, South Western Sydney Local Health District. Australia.

Katterl, R., Anikeeva, O., Butler, C., Brown, L., Smith, B., and Bywood, P. 2012. Potentially avoidable hospitalisations in Australia: Causes for hospitalisations and primary health care interventions. *PHC RIS Policy Issue Review*. Adelaide: Primary Health Care Research & Information Service.

Lawn, S., Zabeen, S., Smith, D., Wilson, E., Miller, C., Battersby, M., and Masman, K. 2018. Managing chronic conditions care across primary care and hospital systems: lessons from an Australian hospital avoidance risk program using the Flinders Chronic Condition Management Program. *Australian Health Review* 42: 542-549.

Longman, J.M., Passey, M.E., Ewald, D.P., Rix, E., and Morgan, G.G. 2015. Admissions for chronic ambulatory care sensitive conditions – a useful measure of potentially preventable admission? *BMC Health Services Research* 15:472.

Longman, J.M., Rix, E., Johnston, J.J., and Passey, M.E. 2018. Ambulatory care sensitive chronic conditions: what can we learn from patients about the role of primary health care in preventing admissions? *Australian Journal of Primary Health* 24: 304-310.

Manski-Nankervis, J., Furler, J., Audehm, R., Blackberry, I., and Young, D. 2015. Potentially preventable hospitalisations: are they a useful marker of access to and experience of care in general practice among people with type 2 diabetes? *Australian Journal of Primary Health* 21:214-220.

- Mercier, G., Georgescu, V., and Bousquet, J. 2015. Geographic variations in potentially avoidable hospitalizations in France. *Health Affairs* 34(5): 836-843.
- Moorin, R.E., Youens, D., Preen, D.B., Harris, M and Wright, C.M. 2019. Association between continuity of provider-adjusted regularity of general practitioner contact and unplanned diabetes-related hospitalisation: a data linkage study in New South Wales, Australia, using the 45 and up study cohort. *BMJ Open* 9:e027158.
- Muenchberger, H. and Kendall, E. 2010. Predictors of preventable hospitalisation in chronic disease: Priorities for change. *Journal of Public Health Policy*. 32 (2): 150-163.
- National Health Performance Authority (NHPA). 2015. *Healthy Communities: Potentially preventable hospitalisations in 2013–14, Technical Supplement*. Sydney: NHPA.
- National Health Performance Authority (NHPA). 2017. *Healthy Communities: Potentially Preventable Hospitalisations in 2015-2016*. Sydney: NHPA.
- Page, A., Ambrose, S., Glover, J., and Hetzel, D. 2007. *Atlas of Avoidable Hospitalisations in Australia: ambulatory care – sensitive conditions*. Adelaide: PHIDU. University of Adelaide.
- Passey, M.E., Longman, J.M., Johnston, J.J., Jorm, L., Ewald, D., Morgan, G.G., Rolfe, M., Chalker, B. 2015. Diagnosing Potentially Preventable Hospitalisations (DaPPHne): protocol for a mixed-methods data-linkage study. *BMJ Open* 2015;5: e009879. doi:10.1136/ bmjopen-2015-00987.
- Passey, M.E. 2018. DaPPHne (Diagnosing Potentially Preventable Hospitalisations): Understanding unplanned hospital admissions for chronic conditions. <http://ruralhealth.org.au/6rrhss/sites/default/files/2018-04/1150%20Wed%20Cbr%20Megan%20Passey.pdf> (accessed 26/06/2019).
- Public Health Information Development Unit. 2015. *The socioeconomic gradient and chronic conditions in Australia: Results from the 2011-13 Australian Health Survey*. Torrens University Australia. Adelaide. PHIDU.
- Public Health Information Development Unit. 2018. *Potentially preventable hospitalisations in Australia: variations by sociodemographic characteristics and geographical areas, with a focus on Aboriginal and Torres Strait Islander people, 2012/13 to 2014/15*. Torrens University Australia. Adelaide: PHIDU.
- Purdy S., Griffin, T., Salisbury, C., and Sharp, D. 2009. Ambulatory care sensitive conditions: terminology and disease coding need to be more specific to aid policy makers and clinicians. *Public Health* 123:169-173.
- Purdy, S. and Huntley, A. 2013. Predicting and preventing avoidable hospital admissions: A review. *J R Coll Physicians Edinb*, 43: 340-4.
- Pollmans, J., Romano, P.S., Weyermann, M., Geraedts, M., and Drosler, S.E. 2018. Impact of disease prevalence adjustment on hospitalisation rates for chronic ambulatory care-sensitive conditions in Germany. *Health services research*. 53(2): 1180- 1202.
- Rosano, A., Loha, C.A., Falvo, R., van der Zee, J., Ricciardi, W., Guasticchi, G., and de Belvis, A.G. 2012. The relationship between avoidable hospitalization and accessibility to primary care: a systematic review. *European Journal of Public Health* 23(3): 356-360.
- Solberg, L.I. 2015. Preventable hospital admissions: are they? *Family Practice* 32(3):245-246.

Sundmacher, L., Fischbach, D., Schuettig, W., Naumann, U.A, and Faisst, C. 2015. Which hospitalisations are ambulatory care-sensitive, to what degree and how could the rates be reduced? Results of a group consensus study in Germany. *Health Policy* 119: 1415-1423.

Swanson, J.O., Vogt, V., Sundmacher, L., Hagen, T.P., and Moger, T.A. 2018. Continuity of care and its effect on readmission for COPD patients: A comparative study in Norway and Germany. *Health Policy* 122: 737-745.

Tran, B., Falster, M.O., Douglas, K., Blyth, F., and Jorm, L.R. 2014. Health behaviours and potentially preventable hospitalisation: A prospective study of older Australian adults. *PLOS One* 9:4. E931111.

van Loenen, T., van den Berg, M.J., Westert, G.P. and Faber, M.J. 2014. Organizational aspects of primary care to avoidable hospitalization: a systematic review. *Family Practice* 31(5): 502-516.

Wolters, R.J., Braspenning, J.C.C, and Wensing, M. 2017. Impact of primary care on hospital admission rates for diabetes patient: A systematic review. *Diabetes Research and Clinical Practice* 129: 182-196.

Wensing, M., Kaufmann, P.K., Szecsenyi, J., Stock, C., and Laux, G. 2018. Effects of a program to strengthen general practice care on hospitalisation rates: a comparative observational study, *Scandinavian Journal of Primary Health Care* 36(2): 109-114.

Walker, R. L., Ghali, W.A., Chen, G., Khalsa, T.K., Mangat, B.K., Campbell, N.R., Dixon, E., Rabli, D., Jette, N., Dhanoa, R., and Quan, H. 2017. ACSC Indicator: testing the reliability for hypertension. *BMC Medical Informatics and Decision Making* 17:90.

World Health Organization (WHO). 2016 Assessing health services delivery performance with hospitalisations for ambulatory care sensitive conditions. Working Document. Health Services Delivery Programme, Division of Health Systems and Public Health.

Yap, M., Kok, M. Nanda, S., Vickery, A., and Whyatt, D. 2018. Hospital admissions and emergency department presentations for dental conditions indicate access too hospital, rather than poor access to dental care in the community. *Australian Journal of Primary Health* 24:74-81.

Zhao, Y, Thomas, S.L., Guthridge, S.L., and Wakeman, J. 2014. Better health outcomes at lower costs: the benefits of primary care utilisation for chronic disease management in remote Indigenous communities in Australia's Northern Territory. *BMC Health Services Research* 14:463.

Zhang, J., Donald, M., Baxter, K.A, Ware, R.S., BurrIDGE, L., Russell, A.W, and Jackson C.L. 2015. Impact of an integrated model of care on potentially preventable hospitalisations for people with Type 2 diabetes mellitus. *Diabetic Medicine*: 872-880.