

Population health profile of the Illawarra

Division of General Practice: supplement

Population Profile Series: No. 15a

PHIDU

March 2007



PHIDU



Australian Government

Australian Institute of
Health and Welfare



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA

© Commonwealth of Australia 2007

This work may be reproduced and used subject to acknowledgement of the source of any material so reproduced.

National Library of Australia Cataloguing in Publication entry

Population health profile of the Illawarra Division of General Practice: supplement.

ISBN 9 78073089 6159 (web).

1. Public health - New South Wales - Illawarra - Statistics.
2. Health status indicators - New South Wales - Illawarra - Statistics. 3. Health service areas - New South Wales - Illawarra. 4. Illawarra (N.S.W.) - Statistics, Medical. I. Public Health Information Development Unit (Australia). (Series : Population profile series ; no. 15a).

362.1099446

ISSN 1833-0452 Population Profile Series

Public Health Information Development Unit, The University of Adelaide *A Collaborating Unit of the Australian Institute of Health and Welfare*

This profile was produced by PHIDU, the Public Health Information Development Unit at The University of Adelaide, South Australia. The work was funded under a grant from the Australian Government Department of Health and Ageing. The views expressed in this profile are solely those of the authors and should not be attributed to the Department of Health and Ageing or the Minister for Health and Ageing.

Interpretation of differences between data in this profile and similar data from other sources needs to be undertaken with care, as such differences may be due to the use of different methodology to produce the data.

Suggested citation:

PHIDU. (2007) *Population health profile of the Illawarra Division of General Practice: supplement*. Population Profile Series: No. 15a. Public Health Information Development Unit (PHIDU), Adelaide.

Enquiries about or comments on this publication should be addressed to:

PHIDU, The University of Adelaide, South Australia 5005
Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au

This publication, the maps and supporting data, together with other publications on population health, are available from the PHIDU website (www.publichealth.gov.au).

Published by Public Health Information Development Unit, The University of Adelaide

Contributors: Anthea Page, Sarah Ambrose, Kristin Leahy and John Glover

Population health profile of the Illawarra Division of General Practice: supplement

This profile is a supplement to the *Population health profile of the Illawarra Division of General Practice*, dated November 2005, available from www.publichealth.gov.au. This supplement includes an update of the population of the Illawarra Division of General Practice, as well as additional indicators and aspects of the Division's socioeconomic status, use of GP services and health. The contents are:


- Population [updated to June 2005]
- Additional socio-demographic indicators
- Unreferred attendances – patient flow/ GP catchment
- Additional prevalence estimates: chronic diseases and risk factors combined
- Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions
- Avoidable mortality

For further information on the way Division totals in this report have been estimated, please refer to the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Population

The Illawarra Division had an Estimated Resident Population of 275,883 at 30 June 2005.

Figure 1: Annual population change, Illawarra DGP, country New South Wales, New South Wales and Australia, 1991 to 1996, 1996 to 2001 and 2001 to 2005




Over the five years from 1991 to 1996, the Division's population increased by 0.9% on average each year, equal to the rate in country New South Wales and lower than in New South Wales (1.0%) and Australia (1.3%). From 1996 to 2001, the annual percentage increase was 1.2%, higher than for country New South Wales (1.1%), but lower than New South Wales or Australia (both 1.3%). The growth rate declined to 0.4% per year from 2001 to 2005, below the annual increases for country New South Wales and New South Wales (0.7% and 0.6%, respectively).

Table 1: Population by age, Illawarra DGP and Australia, 2005

Age group (years)	Illawarra DGP		Australia	
	No.	%	No.	%
0-14	54,407	19.7	3,978,221	19.6
15-24	38,261	13.9	2,819,834	13.9
25-44	76,578	27.8	5,878,107	28.9
45-64	65,939	23.9	4,984,446	24.5
65-74	21,756	7.9	1,398,831	6.9
75-84	14,755	5.3	954,143	4.7
85+	4,187	1.5	315,027	1.5
Total	275,883	100.0	20,328,609	100.0

As shown in the accompanying table and the age-sex pyramid below, Illawarra DGP had similar proportions of its population in the two youngest age groups and slightly lower proportions in the 25 to 44 years (27.8%) and 45 to 64 years (23.9%) compared to Australia (with 28.9% and 24.5%, respectively). The 65 to 74 years (7.9%) and 75 to 84 years age groups (5.3%) had higher proportions compared to Australia (6.9% and 4.7%).


Figure 2: Population in Illawarra DGP and Australia, by age and sex, 2005



The age distribution of the Division's population is similar to that for Australia. The only notable differences are:

- from 25 to 39 years – generally lower proportions of both females and males;
- from 45 to 59 years – slightly lower proportions of males and, females; and
- at older ages – marginally higher proportions of males aged 65 to 84 years and females aged 60 to 84 years.

Figure 3: Population projections for Illawarra DGP, by age and sex, 2005 and 2020




The population projections for the Division show a number of changes in age distribution, with the 2020 population projected to have:

- at younger ages – much lower proportions of males and females aged 0 to 34 years (only marginally lower at ages 25 to 29 years) and lower proportions at ages 40 to 49 years; and
- from ages 50 years and onwards – higher proportions of males and females (most pronounced at ages 60 to 74 years).

Additional socio-demographic indicators

Please refer to the earlier *Population health profile of the Illawarra Division of General Practice*, dated November 2005, available from www.publichealth.gov.au, for other socio-demographic indicators.

Figure 4: Index of Relative Socio-Economic Disadvantage, Illawarra DGP, 2001



One of four socioeconomic indexes for areas produced at the 2001 ABS Census is the Index of Relative Socio-Economic Disadvantage.

The Illawarra DGP has an index score of 981, below the score for Australia of 1000: this score varies widely across the Division, from 825 in the most disadvantaged areas to 1094 in the least disadvantaged areas.


Note: each 'quintile' comprises approximately 20% of the population of the Division.

A new indicator, produced for the first time at the 2001 ABS Census, shows the number of jobless families with children under 15 years of age. There were fewer jobless families in the Illawarra DGP (18.3%), compared to country New South Wales as a whole (20.9%) (Figure 5, Table 2).

With the introduction of the 30% rebate for private health insurance premiums, there was a once-off registration process, providing information of the postcode and residence of those who had such insurance (these data are not available at this area level for later dates). In 2001, the proportion of the population with private health insurance (45.0%), was consistent with the level in country New South Wales (44.9%) (Figure 5, Table 2).

Figure 5: Socio-demographic indicators, Illawarra DGP, country New South Wales, New South Wales and Australia, 2001

Jobless families with children under 15 years old



Private health insurance, 30 June





Table 2: Socio-demographic indicators, Illawarra DGP, country New South Wales, New South Wales and Australia, 2001

Indicator	Illawarra DGP		Country NSW		New South Wales		Australia	
	No.	%	No.	%	No.	%	No.	%
Jobless families with children under 15 years old	5,284	18.3	54,883	20.9	121,409	17.6	357,563	17.4
Private health insurance (30 June)	115,825	45.0	1,061,580	44.9	3,062,382	48.2	8,671,106	46.0


Details of the distribution of jobless families and of the population covered by private health insurance are shown by Statistical Local Area (SLA) in Maps 1 and 2, respectively.

Map 1: Jobless families with children under 15 years of age by SLA, Illawarra DGP, 2001



[#] data were not mapped: see 'Mapping' note under Methods

Map 2: People covered by private health insurance by SLA, Illawarra DGP, 30 June 2001



[#] data were not mapped: see 'Mapping' note under Methods

GP services to residents of the Illawarra DGP

The following tables include information, purchased from Medicare Australia, of the movement of patients and GPs between Divisions. Note that the data only include unreferred attendances recorded under Medicare: unreferred attendances not included are those for which the cost is met by the Department of Veterans' Affairs or a compensation scheme; or are provided by salaried medical officers in hospitals, community health services or Aboriginal Medical Services, and which are not billed to Medicare. At any attendance, one or more services may have been provided.

The majority (94.5%) of all unreferred attendances to residents of Illawarra DGP were provided in the Division (i.e. by a GP with a provider number in the Division): this represented 1,468,635 GP unreferred attendances (Table 3). The remainder of unreferred attendances to residents were provided by GPs with a provider number in surrounding and other Divisions.

Table 3: Patient flow – People living¹ in Illawarra DGP by Division where attendance occurred², 2003/04

Division		Unreferred attendances	
Number	Name	No.	% ³
216	Illawarra DGP	1,468,635	94.5
214	Sutherland DGP	11,654	0.8
202	Eastern Sydney DGP	8,050	0.5
209	St George DGP	6,479	0.4
201	Central Sydney DGP	6,096	0.4
Other	..	52,741	3.4
Total	..	1,553,655	100.0

¹ Based on address in Medicare records

² Division of GP based on provider number

³ Proportion of all unreferred attendances of patients with an address in Division 216 by Division in which attendance occurred

The majority (95.1%) of unreferred attendances provided by GPs with a provider number in Illawarra DGP were also to people living in the Division (i.e. their Medicare address was in the Division) (Table 4). The remainder of unreferred attendances to residents were provided by GPs with a provider number in surrounding and other Divisions.

Table 4: GP catchment – Unreferred attendances provided by GPs¹ in Illawarra DGP by Division of patient address², 2003/04

Division		Unreferred attendances	
Number	Name	No.	% ³
216	Illawarra DGP	1,468,635	95.1
220	Shoalhaven DGP	9,071	0.6
215	Macarthur DGP	6,309	0.4
235	Southern Highlands DGP	4,913	0.3
214	Sutherland DGP	4,658	0.3
221	South East NSW DGP	4,339	0.3
Other	..	46,123	3.0
Total	..	1,544,048	100.0

¹ Division of GP based on provider number

² Based on address in Medicare records

³ Proportion of all unreferred attendances to GPs with a provider number in Division 216 by Division of patient address

Additional prevalence estimates: chronic diseases and risk factors combined

Please refer to the earlier *Population health profile of the Illawarra Division of General Practice*, dated November 2005, available from www.publichealth.gov.au, for the separate prevalence estimates of chronic disease; measures of self-reported health and risk factors. The process by which the estimates have been made, and details of their limitations, are also described in the 'Notes on the data' section of this earlier profile.

In this section two estimates, which combine the prevalence of selected chronic diseases with a risk factor, are shown for the Division. The measures are of people who *had asthma and were smokers*, and people who *had type 2 diabetes and were overweight or obese*: note that the estimates have been predicted from self-reported data, and are not based on clinical records or physical measures.

It is estimated that there were relatively more people in Illawarra DGP who had asthma and were smokers, compared to Australia as a whole, but fewer than in country New South Wales (Figure 6, Table 5): that is, the prevalence rates per 1,000 population were higher than the national rates. There were higher rates in Illawarra DGP of people who had type 2 diabetes and were overweight or obese, compared to country New South Wales or Australia.

Figure 6: Estimates of selected chronic diseases and risk factors, Illawarra DGP, country New South Wales and Australia, 2001



Table 5: Estimates of selected chronic diseases and risk factors, Illawarra DGP, country New South Wales, New South Wales and Australia, 2001

Variable	Illawarra DGP		Country NSW		New South Wales		Australia	
	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ¹
Had asthma and smoked ³	5,800	22.5	54,344	24.7	126,542	19.7	397,734	20.8
Had type 2 diabetes & were overweight/ obese ⁴	4,707	17.8	40,784	15.5	100,235	15.7	283,176	15.2

¹ No. is a weighted estimate of the number of people in Illawarra DGP reporting these chronic conditions/ with these risk factors and is derived from synthetic predictions from the 2001 NHS

² Rate is the indirectly age-standardised rate per 1,000 population

³ Population aged 18 years and over

⁴ Population aged 15 years and over

Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions

The rationale underlying the concept of avoidable hospitalisations is that timely and effective care of certain conditions, delivered in a primary care setting, can reduce the risk of hospitalisation. Admissions to hospital for these ambulatory care sensitive (ACS) conditions can be avoided in three ways. Firstly, for conditions that are usually preventable through immunisation or nutritional intervention, disease can be prevented almost entirely. Secondly, diseases or conditions that can lead to rapid onset problems, such as dehydration and gastroenteritis, can be treated. Thirdly, chronic conditions, such as congestive heart failure, can be managed to prevent or reduce the severity of acute flare-ups to avoid hospitalisation.

This measure does not include other aspects of avoidable morbidity, namely potentially preventable hospitalisations (hospitalisations resulting from diseases preventable through population based health promotion strategies, e.g. alcohol-related conditions; and most cases of lung cancer) and hospitalisations avoidable through injury prevention (e.g. road traffic accidents).

For information on the ambulatory care sensitive conditions and ICD codes included in the analysis in this section, please refer to the *Atlas of Avoidable Hospitalisations in Australia: ambulatory care-sensitive conditions*, available from www.publichealth.gov.au.

In 2001 to 2002, the 7,414 admissions from ambulatory care sensitive (ACS) conditions accounted for 8.5% of all admissions in the Illawarra DGP (Table 6, Figure 7): this was consistent with the levels for both New South Wales (8.6%) and Australia (8.7%).


Table 6: Avoidable¹ and unavoidable hospitalisations, Illawarra DGP, New South Wales, and Australia, 2001/02

Category	Illawarra DGP			New South Wales			Australia		
	No.	Rate ²	%	No.	Rate ²	%	No.	Rate ²	%
Avoidable ¹	7,414	2,673.6	8.5	170,066	2,543.8	8.6	552,786	2,847.5	8.7
Unavoidable	79,849	29,152.0	91.5	1,810,901	27,255.3	91.4	5,818,199	29,970.7	91.3
Total	87,263	31,824.5	100.0	1,980,967	29,798.8	100.0	6,370,985	32,818.2	100.0

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

Figure 7: Avoidable hospitalisations¹, Illawarra DGP, New South Wales and Australia, 2001/02




The rate of avoidable hospitalisations in Illawarra DGP, 2,673.6 admissions per 100,000 population, is above that in New South Wales (a rate of 2,543.8) but below that in Australia (2,847.5).

¹ Admissions resulting from ACS conditions

Diabetes complications, chronic obstructive pulmonary disease, dental conditions and congestive heart failure were the four conditions with the highest rates of avoidable hospitalisations in the Illawarra DGP (Figure 8, Table 7).

Table 7 shows the number, rate and proportion of avoidable hospitalisations, for the individual ACS conditions, as well as the vaccine-preventable; acute; and chronic sub-categories. Almost two-thirds of avoidable hospitalisations are attributable to chronic health conditions. The predominance of hospitalisations for chronic conditions in this period can be primarily attributed to the large number of admissions for diabetes complications. Dental conditions; and convulsions and epilepsy have the highest rates of avoidable hospitalisations for the acute conditions.

Figure 8: Avoidable hospitalisations¹ by condition, Illawarra DGP and New South Wales, 2001/02



¹ Admissions resulting from ACS conditions: excludes nutritional deficiencies as less than ten admissions

Table 7: Avoidable hospitalisations¹ by condition, Illawarra DGP, New South Wales and Australia, 2001/02

Sub-category/ condition	Illawarra DGP		New South Wales		Australia	
	No.	Rate ²	No.	Rate ²	No.	Rate ²
Vaccine-preventable	191	69.6	5,630	84.5	16,573	85.4
Influenza and pneumonia	155	56.3	4,280	64.1	13,021	67.1
Other vaccine preventable	36	13.3	1,350	20.4	3,552	18.3
Chronic³	4,816	1,712.7	106,803	1,587.0	352,545	1,816
Diabetes complications	1,680	593.3	34,975	519.5	141,345	728.1
Iron deficiency anaemia	175	63.4	4,494	67.0	16,451	84.7
Hypertension	81	29.2	2,398	35.7	6,354	32.7
Congestive heart failure	701	252.9	14,270	209.7	42,447	218.6
Angina	679	241.8	16,987	251.8	49,963	257.4
Chronic obstructive pulmonary disease	974	339.0	19,359	285.6	54,853	282.6
Asthma	526	193.1	14,289	216.8	41,009	211.3
Acute	2,681	988.0	62,543	946.0	200,913	1,035
Dehydration and gastroenteritis	476	175.0	11,725	176.4	37,766	194.5
Convulsions and epilepsy	484	178.5	11,093	168.1	31,137	160.4
Ear, nose and throat infections	302	110.8	10,615	161.1	32,075	165.2
Dental conditions	789	291.5	11,196	170.3	43,667	224.9
Perforated/bleeding ulcer	77	27.7	1,830	27.1	5,795	29.9
Ruptured appendix	72	26.7	1,212	18.5	3,866	19.9
Pyelonephritis	59	21.9	2,038	31.0	7,386	38.0
Pelvic inflammatory disease	83	31.9	2,134	32.7	6,547	33.7
Cellulitis	294	107.8	9,451	142.0	28,204	145.3
Gangrene	45	16.2	1,249	18.6	4,470	23.0
Total avoidable hospitalisations⁴	7,414	2,673.6	170,066	2,543.8	552,786	2,847.5

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

³ Excludes nutritional deficiencies as less than ten admissions

⁴ Sub-category and condition numbers and rates do not add to the reported total avoidable admissions: five conditions (influenza & pneumonia, other vaccine preventable, diabetes complications, ruptured appendix and gangrene) are counted in 'any diagnosis', so may be included in more than one condition group

Avoidable mortality

Avoidable and amenable mortality comprises those causes of death that are potentially avoidable at the present time, given available knowledge about social and economic policy impacts, health behaviours, and health care (the latter relating to the subset of amenable causes).

For information on the avoidable and amenable mortality conditions and ICD codes included in the analysis in this section, please refer to the *Australian and New Zealand Atlas of Avoidable Mortality*, available from www.publichealth.gov.au.

Almost three quarters (71.3%) of all deaths in Illawarra DGP at ages 0 to 74 years over the period 1997 to 2001 are considered to be avoidable, consistent with that for country New South Wales (71.6%) (Table 8). Deaths amenable to health care (amenable mortality, a subset of avoidable mortality) accounted for 28.2% of all deaths at ages 0 to 74 years in Illawarra DGP, compared to 28.3% in country New South Wales.

Table 8: Avoidable and unavoidable mortality (0 to 74 years) by area, Illawarra DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Mortality category	Illawarra DGP		Country NSW		New South Wales		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable	2,904	219.2	29,442	234.3	66,151	213.6	189,845	211.8
% of total	71.3	..	71.6	..	71.4	..	71.5	..
(Amenable)	(1,151)	(86.3)	(11,638)	(91.2)	(26,374)	(85.0)	(76,249)	(85.1)
(% of total)	(28.2)	(..)	(28.3)	(..)	(28.5)	(..)	(28.7)	(..)
Unavoidable	1,171	87.9	11,700	92.1	26,468	85.3	75,582	84.3
% of total	28.2	..	28.4	..	28.6	..	28.5	..
Total mortality	4,075	307.1	41,142	326.4	92,619	299.0	265,427	296.1
%	100.0	..	100.0	..	100.0	..	100.0	..

¹ Rate is the indirectly age-standardised rate per 100,000 population

Rates of avoidable mortality were higher for males than for females in each of the comparator areas. Illawarra DGP's rate of avoidable mortality for males was 287.7 deaths per 100,000 males, notably higher than the rate of 149.7 for females. Similarly, the rate of amenable mortality for males in the Division was higher, 99.7, compared to 72.8 for females, a rate ratio of 1.37 (Figure 9, Table 9).

Figure 9: Avoidable and amenable mortality by sex (0 to 74 years), Illawarra DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Note: the different scales




Table 9: Avoidable and amenable mortality (0 to 74 years) by sex, Illawarra DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Mortality category and sex	Illawarra DGP		Country NSW		New South Wales		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
Males	1,910	287.7	19,569	308.5	43,074	276.8	123,026	272.6
Females	994	149.7	9,873	159.1	23,077	149.6	66,819	150.1
Total	2,904	219.2	29,442	234.3	66,151	213.6	189,845	211.8
Rate ratio-M:F²	..	1.92**	..	1.94**	..	1.85**	..	1.82**
Amenable								
Males	668	99.7	6,743	103.6	14,811	94.8	42,568	94.3
Females	483	72.8	4,895	78.6	11,562	74.9	33,681	75.7
Total	1,151	86.3	11,638	91.2	26,374	85.0	76,249	85.1
Rate ratio-M:F²	..	1.37**	..	1.32**	..	1.27**	..	1.25**

¹ Rate is the indirectly age-standardised rate per 100,000 population

² Rate ratio (M:F) is the ratio of male to female rates; rate ratios differing significantly from 1.0 are shown with * p <0.05; ** p <0.01

Another way of measuring premature mortality is to calculate the number of years of life lost (YLL)¹, which takes into account the years a person could have expected to live at each age of death based on the average life expectancy at that age.

The numbers of YLL for Illawarra DGP, country New South Wales, New South Wales and Australia over the period of analysis are shown in Table 10 by mortality category. However, given the substantial variation in the populations of these areas, a comparison of the proportion of YLL for each area is also shown.

YLL from avoidable mortality accounted for 71.7% of total YLL (0 to 74 years) for Illawarra DGP, consistent with 71.8% for country New South Wales. Similarly, the proportion of YLL from unavoidable mortality for Illawarra DGP (28.3%) was consistent with that for country NSW (28.2%).

Table 10: Years of life lost from avoidable mortality (0 to 74 years), Illawarra DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Mortality category	Illawarra DGP		Country NSW		New South Wales		Australia	
	No.	% of total	No.	% of total	No.	% of total	No.	% of total
Avoidable	49,376	71.7	502,860	71.8	1,147,183	71.8	3,327,375	71.9
(Amenable)	(19,058)	(27.7)	(192,960)	(27.6)	(444,143)	(27.8)	(1,298,430)	(28.0)
Unavoidable	19,474	28.3	197,182	28.2	451,496	28.2	1,303,289	28.1
Total	68,850	100.0	700,042	100.0	1,598,679	100.0	4,630,664	100.0

¹ Years of life lost were calculated using the remaining life expectancy method (this provides an estimate of the average time a person would have lived had he or she not died prematurely). The reference life table was the Coale and Demeny Model Life Table West level 26 female (for both males and females), with the YLL discounted to net present value at a rate of 3 per cent per year.

In each of the areas in Table 11, the majority of avoidable mortality at ages 0 to 74 years occurred in the 65 to 74 year age group (Table 11), with 1,416.2 deaths per 100,000 population in Illawarra Division. The 45 to 64 year age group accounted for the next highest rate of avoidable death in all of the comparators, with a rate 328.3 in Illawarra Division.

Table 11: Avoidable and amenable mortality by age, Illawarra DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Mortality category and age (years)	Illawarra DGP		Country NSW		New South Wales		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
0-14	58	20.9	738	29.0	1,836	27.5	5,669	28.8
15-24	79	41.7	938	62.6	2,241	50.9	7,045	52.8
25-44	340	87.1	3,317	99.6	8,119	82.9	24,356	83.9
45-64	976	328.3	9,755	343.5	22,358	311.1	64,282	304.9
65-74	1,451	1,416.2	14,694	1464.0	31,597	1,375.8	88,493	1,358.1
Total	2,904	219.2	29,442	234.3	66,151	213.6	189,845	211.8
Amenable								
0-24	59	12.5	645	15.5	1,658	14.8	5,083	15.4
25-44	87	22.2	784	23.0	1,878	19.2	5,946	20.5
45-64	396	132.7	4,060	142.9	9,444	131.4	27,464	130.3
65-74	611	595.6	6,148	613.7	13,394	582.9	37,756	579.4
Total	1,151	86.3	11,638	91.2	26,374	85.0	76,249	85.1

¹ Rate is the indirectly age-standardised rate per 100,000 population

Table 12 shows the number and age-standardised death rate by selected major condition group and selected causes included in the avoidable mortality classification.

The highest rates of avoidable mortality for the selected major condition groups in the Illawarra DGP were for cardiovascular diseases, with a rate of 77.5 deaths per 100,000 population, and cancer, 72.1 deaths per 100,000 population (Table 12, Figure 10). For the selected causes within the condition groups, the two major causes of avoidable mortality were ischaemic heart disease and lung cancer, with rates of 60.1 per 100,000 population and 27.4 per 100,000, respectively.


Table 12: Avoidable mortality (0 to 74 years) by major condition group and selected cause, Illawarra DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Condition group/ selected cause	Illawarra DGP		Country NSW		New South Wales		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Cancer	964	72.1	9,239	71.9	21,158	68.1	62,338	69.5
Colorectal cancer	193	14.4	1,936	14.9	4,318	13.9	13,008	14.5
Lung cancer	370	27.4	3,314	25.2	7,297	23.4	21,208	23.7
Cardiovascular diseases	1,048	77.5	10,101	77.0	21,925	70.3	59,945	66.9
Ischaemic heart disease	812	60.1	7,474	57.0	15,935	51.1	43,712	48.8
Cerebrovascular diseases	199	14.7	2,015	15.4	4,656	14.9	12,558	14.0
Respiratory system diseases	158	11.5	2,136	16.0	4,313	13.8	11,612	13.0
Chronic obstructive pulmonary disease	141	10.2	1,966	14.6	3,882	12.4	10,395	11.6
Unintentional injuries	164	13.1	2,027	18.6	4,540	15.0	14,224	15.9
Road traffic injuries	81	6.5	1,279	11.8	2,528	8.4	8,138	9.1
Intentional injuries	196	15.9	1,939	18.1	4,497	14.9	13,891	15.5
Suicide and self inflicted injuries	170	13.8	1,730	16.1	3,941	13.0	12,393	13.8

¹ Rate is the indirectly age-standardised rate per 100,000 population

Rates in the Division for cancers and cardiovascular diseases were generally above those for Australia, but for respiratory system diseases, unintentional injuries and road traffic injuries rates were lower (Figure 10).

Figure 10: Avoidable mortality (0 to 74 years) by major condition group and selected cause, Illawarra DGP, country New South Wales and Australia, 1997 to 2001



Notes on the data

Data sources and limitations

General

References to 'country New South Wales' relate to New South Wales excluding the Sydney Statistical Division.

Data sources

Table 13 details the data sources for the material presented in this profile.

Table 13: Data sources

Section	Source
Population	
Figures 1 and 2; Table 1	Estimated Resident Population, ABS, 30 June for the periods shown
Figure 3	Estimated Resident Population, ABS, 30 June 2005; Population Projections, ABS, 30 June 2020 (unpublished) ¹
Additional socio-demographic indicators	
Figure 4	ABS SEIFA package, Census 2001
Table 2; Figure 5; Map 1	Jobless families, ABS, 2001 (unpublished)
Table 2; Figure 5; Map 2	Private health insurance, from Hansard
GP services – patient flow/ GP catchment	
Tables 3 and 4	Medicare Australia, 2003/04
Additional prevalence estimates: chronic diseases and risk factors combined	
Figure 6; Table 5	Estimated from 2001 National Health Survey (NHS), ABS (unpublished)
Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions	
Tables 6 and 7; Figures 7 and 8	National Hospital Morbidity Database at Australian Institute of Health & Welfare, 2001/02; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)
Avoidable mortality	
Tables 8, 9, 10, 11 and 12; Figures 9 and 10	ABS Deaths 1997-2001; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)

¹ The projected population at June 2020 is based on the 2002 ERP. As such, it is somewhat dated, and does not take into account more recent demographic trends: it is however the only projection series available at the SLA level for the whole of Australia.

Methods

For background information on the additional prevalence estimates presented in this profile, please refer to the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Please also refer to the November 2005 profile for information on the data converters.

Mapping

In some Divisions the maps may include a very small part of an SLA which has not been allocated any population; or has a population of less than 100 or has less than 1% of the SLAs total population; or there were less than five cases (i.e. jobless families, people with health insurance): these areas are mapped with a pattern.

Statistical geography of the Illawarra DGP

For information on the postcodes in the Division, please refer the Department of Health and Ageing website <http://www.health.gov.au/internet/wcms/publishing.nsf/Content/health-pcd-programs-divisions-divspc.htm>; also included in table format in the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Statistical Local Areas (SLAs) are defined by the Australian Bureau of Statistics to produce areas for the presentation and analysis of data. Statistical Local Areas (SLAs) in this Division are the same as the Local Government Areas (Table 14).

Table 14: SLAs and population in Illawarra DGP, 2005 on 2001 boundaries

SLA code	SLA name	Per cent of the SLA's population in the Division *	Estimate of the SLA's 2005 population in the Division
14400	Kiama	100.0	20,357
16900	Shellharbour	100.0	63,124
18450	Wollongong	100.0	192,402

* Proportions are approximate and are known to be incorrect in some cases, due to errors in the concordance used to allocate CDs to form postal areas

Acknowledgements

Funding for these profiles was provided by the Population Health Division of the Department of Health and Ageing (DoHA).

Further developments and updates

When the re-aligned boundaries are released and DoHA have made known their geographic composition, PHIDU will examine the need to revise and re-publish these profiles (*Population health profile*, dated November 2005, and the *Population health profile: supplement*, dated March 2007).

PHIDU contact details

For general comments, data issues or enquiries re information on the web site, please contact PHIDU:

Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au