

**Population health profile of the
Hawkesbury-Hills
Division of General Practice
(formerly Hawkesbury DGP): supplement**

Population Profile Series: No. 36a

PHIDU

March 2007



© Commonwealth of Australia 2007

This work may be reproduced and used subject to acknowledgement of the source of any material so reproduced.

National Library of Australia Cataloguing in Publication entry

Population health profile of the Hawkesbury-Hills Division of General Practice (formerly Hawkesbury DGP): supplement.

Bibliography.

ISBN 9 78073089 6340 (web).

1. Public health - New South Wales - Hawkesbury - Statistics.
 2. Health status indicators - New South Wales - Hawkesbury - Statistics.
 3. Health service areas - New South Wales - Hawkesbury.
 4. Hawkesbury (N.S.W.) - Statistics, Medical.
- I. Public Health Information Development Unit (Australia).
(Series : Population profile series ; no. 36a).

362.1099442

ISSN 1833-0452 Population Profile Series

Public Health Information Development Unit, The University of Adelaide *A Collaborating Unit of the Australian Institute of Health and Welfare*

This profile was produced by PHIDU, the Public Health Information Development Unit at The University of Adelaide, South Australia. The work was funded under a grant from the Australian Government Department of Health and Ageing. The views expressed in this profile are solely those of the authors and should not be attributed to the Department of Health and Ageing or the Minister for Health and Ageing.

Interpretation of differences between data in this profile and similar data from other sources needs to be undertaken with care, as such differences may be due to the use of different methodology to produce the data.

Suggested citation:

PHIDU. (2007) *Population health profile of the Hawkesbury-Hills Division of General Practice (formerly Hawkesbury DGP): supplement*. Population Profile Series: No. 36a. Public Health Information Development Unit (PHIDU), Adelaide.

Enquiries about or comments on this publication should be addressed to:

PHIDU, The University of Adelaide, South Australia 5005
Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au

This publication, the maps and supporting data, together with other publications on population health, are available from the PHIDU website (www.publichealth.gov.au).

Published by Public Health Information Development Unit, The University of Adelaide

Contributors: Anthea Page, Sarah Ambrose, Kristin Leahy and John Glover

Population health profile of the Hawkesbury-Hills Division of General Practice (formerly Hawkesbury DGP): supplement

This profile is a supplement to the *Population health profile of the Hawkesbury Division of General Practice* (now known as Hawkesbury-Hills DGP), dated November 2005, available from www.publichealth.gov.au. This supplement includes an update of the population of the Hawkesbury-Hills Division of General Practice, as well as additional indicators and aspects of the Division's socioeconomic status, use of GP services and health. The contents are:


- Population [updated to June 2005]
- Additional socio-demographic indicators
- Unreferred attendances – patient flow/ GP catchment
- Additional prevalence estimates: chronic diseases and risk factors combined
- Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions
- Avoidable mortality

For further information on the way Division totals in this report have been estimated, please refer to the 'Notes on the data' section of the *Population health profile, November 2005* (www.publichealth.gov.au).

Population

The Hawkesbury-Hills Division had an Estimated Resident Population of 240,322 at 30 June 2005.

Figure 1: Annual population change, Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 1991 to 1996, 1996 to 2001 and 2001 to 2005




Over the five years from 1991 to 1996, the Division's population increased by 1.5% on average each year, markedly above the level in country New South Wales (0.9%) and New South Wales (1.0%). From 1996 to 2001, the annual percentage increase in the Division was 2.3%, substantially higher than in country New South Wales (1.1%) and New South Wales (1.3%). The growth rate of 2.1% per year from 2001 to 2005 was again substantially higher than in country New South Wales and New South Wales (0.6%).

Table 1: Population by age, Hawkesbury-Hills DGP and Australia, 2005

Age group (years)	Hawkesbury-Hills DGP		Australia	
	No.	%	No.	%
0-14	53,437	22.2	3,978,221	19.6
15-24	35,723	14.9	2,819,834	13.9
25-44	72,562	30.2	5,878,107	28.9
45-64	58,540	24.4	4,984,446	24.5
65-74	11,527	4.8	1,398,831	6.9
75-84	6,560	2.7	954,143	4.7
85+	1,973	0.8	315,027	1.5
Total	240,322	100.0	20,328,609	100.0

As shown in the accompanying table and the age-sex pyramid below (Figure 2), Hawkesbury-Hills DGP had a higher proportion of the population aged 0 to 14 years, 15 to 24 years and 25 to 44 years (22.2%, 14.9% and 30.2%) compared to Australia (19.6%, 13.9% and 28.9%) (Table 1). Conversely, the 65 years and over age groups had lower proportions compared to Australia as a whole; at the oldest these were markedly lower.


Figure 2: Population in Hawkesbury-Hills DGP and Australia, by age and sex, 2005



The age distribution of the Division's population is similar to that for Australia overall. The most notable differences are:

- at the youngest ages – lower proportion of children and young people aged 0 to 24 years;
- from 30 to 44 years – lower proportions of both males and, more noticeably, of females; and
- at older ages – higher proportions of both males and females from 60 years of age.

Figure 3: Population projections for Hawkesbury-Hills DGP, by age and sex, 2005 and 2020




The population projections for the Division show a number of changes in age distribution, with the 2020 population projected to have:

- at younger ages – much lower proportions of males and females aged 0 to 24 years;
- from 30 to 54 years – lower proportions of males (to 44 years) and females; and
- from age 55 years and onwards – higher proportions of males and females (most pronounced at ages 60 to 74 years).

Additional socio-demographic indicators

Please refer to the earlier *Population health profile of the Hawkesbury Division of General Practice* (now known as Hawkesbury-Hills DGP), dated November 2005, available from www.publichealth.gov.au, for other socio-demographic indicators.

Figure 4: Index of Relative Socio-Economic Disadvantage, Hawkesbury-Hills DGP, 2001



One of four socioeconomic indexes for areas produced at the 2001 ABS Census is the Index of Relative Socio-Economic Disadvantage.

The Hawkesbury-Hills DGP has an index score of 1060, above the score for Australia of 1000: this score varies across the Division, from 954 in the most disadvantaged areas to 1127 in the least disadvantaged areas.


Note: each 'quintile' comprises approximately 20% of the population of the Division.

A new indicator, produced for the first time at the 2001 ABS Census, shows the number of jobless families with children under 15 years of age. There were substantially fewer jobless families in the Hawkesbury-Hills DGP (10.0%), compared with country New South Wales (20.9%) (Figure 5, Table 2).

With the introduction of the 30% rebate for private health insurance premiums, there was a once-off registration process, providing information of the postcode and residence of those who had such insurance (these data are not available at this area level for later dates). In 2001, the Division had a markedly higher proportion of the population with private health insurance (56.0%), compared to country New South Wales (44.9%) (Figure 5, Table 2).

Figure 5: Socio-demographic indicators, Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 2001

Jobless families with children under 15 years old



Private health insurance, 30 June






Table 2: Socio-demographic indicators, Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 2001

Indicator	Hawkesbury-Hills DGP		Country NSW		New South Wales		Australia	
	No.	%	No.	%	No.	%	No.	%
Jobless families with children under 15 years old	2,604	10.0	54,883	20.9	121,409	17.6	357,563	17.4
Private health insurance (30 June)	111,338	56.0	1,061,580	44.9	3,062,382	48.2	8,671,106	46.0


Details of the distribution of jobless families and of the population covered by private health insurance are shown by Statistical Local Area (SLA) in Maps 1 and 2, respectively.

Map 1: Jobless families with children under 15 years of age by SLA, Hawkesbury-Hills DGP, 2001



[#] data were not mapped: see 'Mapping' note under Methods

Map 2: People covered by private health insurance by SLA, Hawkesbury-Hills DGP, 30 June 2001



[#] data were not mapped: see 'Mapping' note under Methods

GP services to residents of the former Hawkesbury DGP*

*Note: the data presented for Division 240 in this section, *GP services*, are based on the July 2005 boundaries – reflecting the old Hawkesbury DGP.

The following tables include information, purchased from Medicare Australia, of the movement of patients and GPs between Divisions. Note that the data only include unreferred attendances recorded under Medicare: unreferred attendances not included are those for which the cost is met by the Department of Veterans' Affairs or a compensation scheme; or are provided by salaried medical officers in hospitals, community health services or Aboriginal Medical Services, and which are not billed to Medicare. At any attendance, one or more services may have been provided.

Just over two thirds (67.8%) of all unreferred attendances to residents of Hawkesbury DGP were provided in the Division (ie. by a GP with a provider number in the Division): this represented 347,072 GP unreferred attendances (Table 3). A further 18.3% of unreferred attendances to residents were provided by GPs with a provider number in Western Sydney DGP, with 4.5% provided by GPs in Hornsby Ku-ring-gai Ryde DGP.

Table 3: Patient flow – People living¹ in Hawkesbury DGP by Division where attendance occurred², 2003/04

Division		Unreferred attendances	
Number	Name	No.	% ³
240	Hawkesbury DGP (now part of Hawkesbury-Hills)	347,072	67.8
206	Western Sydney DGP (now WentWest & part Hawkesbury-Hills)	93,899	18.3
212	Hornsby Ku-ring-gai Ryde DGP	23,081	4.5
237	Nepean DGP	21,585	4.2
201	Central Sydney DGP	3,228	0.6
211	Fairfield DGP	3,121	0.6
Other	..	20,129	3.9
Total	..	512,115	100.0

¹ Based on address in Medicare records

² Division of GP based on provider number

³ Proportion of all unreferred attendances of patients with an address in Division 240 by Division in which attendance occurred

Four fifths (80.4%) of unreferred attendances provided by GPs with a provider number in Hawkesbury DGP were to people living in the Division (ie. their Medicare address was in the Division) (Table 4). A further 11.8% of unreferred attendances by GPs in the Division were to people living in Western Sydney DGP, with 2.1% to residents of Nepean DGP.

Table 4: GP catchment – Unreferred attendances provided by GPs¹ in Hawkesbury DGP by Division of patient address², 2003/04

Division		Unreferred attendances	
Number	Name	No.	% ³
240	Hawkesbury DGP (now part of Hawkesbury-Hills)	347,072	80.4
206	Western Sydney DGP (now WentWest & part Hawkesbury-Hills)	50,823	11.8
237	Nepean DGP	9,026	2.1
212	Hornsby Ku-ring-gai Ryde DGP	3,488	0.8
238	Blue Mountains DGP	2,104	0.5
Other	..	18,966	4.4
Total	..	431,479	100.0

¹ Division of GP based on provider number

² Based on address in Medicare records

³ Proportion of all unreferred attendances to GPs with a provider number in Division 240 by Division of patient address

Additional prevalence estimates: chronic diseases and risk factors combined

Please refer to the earlier *Population health profile of the Hawkesbury Division of General Practice* (now known as Hawkesbury-Hills DGP), dated November 2005, available from www.publichealth.gov.au, for the separate prevalence estimates of chronic disease; measures of self-reported health and risk factors. The process by which the estimates have been made, and details of their limitations, are also described in the 'Notes on the data' section of this earlier profile.

In this section two estimates, which combine the prevalence of selected chronic diseases with a risk factor, are shown for the Division. The measures are of people who *had asthma and were smokers*, and people who *had type 2 diabetes and were overweight or obese*: note that the estimates have been predicted from self-reported data, and are not based on clinical records or physical measures.

It is estimated that there were relatively fewer people in Hawkesbury-Hills DGP who had asthma and were smokers, compared to Australia as a whole and (in particular) country New South Wales (Figure 6, Table 5): that is, the prevalence rates per 1,000 population were lower. Similarly, there were fewer people in Hawkesbury-Hills DGP who had type 2 diabetes and were overweight or obese, compared to Australia and country New South Wales.

Figure 6: Estimates of selected chronic diseases and risk factors, Hawkesbury-Hills DGP, country New South Wales and Australia, 2001



Table 5: Estimates of selected chronic diseases and risk factors, Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 2001

Variable	Hawkesbury-Hills DGP		Country NSW		New South Wales		Australia	
	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ¹
Had asthma and smoked ³	3,782	17.1	54,344	24.7	126,542	19.7	397,734	20.8
Had type 2 diabetes & were overweight/ obese ⁴	2,384	13.4	40,784	15.5	100,235	15.7	283,176	15.2

¹ No. is a weighted estimate of the number of people in Hawkesbury-Hills DGP reporting these chronic conditions/ with these risk factors and is derived from synthetic predictions from the 2001 NHS

² Rate is the indirectly age-standardised rate per 1,000 population

³ Population aged 18 years and over

⁴ Population aged 15 years and over

Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions, former Hawkesbury DGP*

The rationale underlying the concept of avoidable hospitalisations is that timely and effective care of certain conditions, delivered in a primary care setting, can reduce the risk of hospitalisation. Admissions to hospital for these ambulatory care sensitive (ACS) conditions can be avoided in three ways. Firstly, for conditions that are usually preventable through immunisation or nutritional intervention, disease can be prevented almost entirely. Secondly, diseases or conditions that can lead to rapid onset problems, such as dehydration and gastroenteritis, can be treated. Thirdly, chronic conditions, such as congestive heart failure, can be managed to prevent or reduce the severity of acute flare-ups to avoid hospitalisation.

This measure does not include other aspects of avoidable morbidity, namely potentially preventable hospitalisations (hospitalisations resulting from diseases preventable through population based health promotion strategies, e.g. alcohol-related conditions; and most cases of lung cancer) and hospitalisations avoidable through injury prevention (e.g. road traffic accidents).

For information on the ambulatory care sensitive conditions and ICD codes included in the analysis in this section, please refer to the *Atlas of Avoidable Hospitalisations in Australia: ambulatory care-sensitive conditions*, available from www.publichealth.gov.au.

***Note: the data presented for Division 240 in this section, *Avoidable hospitalisations*, are based on the July 2005 boundaries – reflecting the old Hawkesbury DGP.**

In 2001 to 2002, the 1,981 admissions from ambulatory care sensitive (ACS) conditions accounted for 8.2% of all admissions in the Hawkesbury DGP (Table 6, Figure 7), slightly lower than the levels in both New South Wales (8.6%) and Australia (8.7%).


Table 6: Avoidable¹ and unavoidable hospitalisations, Hawkesbury DGP, New South Wales, and Australia, 2001/02

Category	Hawkesbury DGP			New South Wales			Australia		
	No.	Rate ²	%	No.	Rate ²	%	No.	Rate ²	%
Avoidable ¹	1,981	2,468.4	8.2	170,066	2,543.8	8.6	552,786	2,847.5	8.7
Unavoidable	22,198	26,168.5	91.8	1,810,901	27,255.3	91.4	5,818,199	29,970.7	91.3
Total	24,179	28,637.9	100.0	1,980,967	29,798.8	100.0	6,370,985	32,818.2	100.0

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

Figure 7: Avoidable hospitalisations¹, Hawkesbury DGP, New South Wales and Australia, 2001/02




The rate of avoidable hospitalisations in Hawkesbury DGP is lower, a rate of 2,468.4 admissions per 100,000 population, compared to both New South Wales (a rate of 2,543.8), and Australia (2,847.5).

¹ Admissions resulting from ACS conditions

Diabetes complications, chronic obstructive pulmonary disease, angina and congestive heart failure were the four conditions with the highest rates of avoidable hospitalisations in the Hawkesbury DGP (Figure 8, Table 7).

Table 7 shows the number, rate and proportion of avoidable hospitalisations, for the individual ACS conditions, as well as the vaccine-preventable; acute; and chronic sub-categories. Almost two-thirds of avoidable hospitalisations are attributable to chronic health conditions. The predominance of hospitalisations for chronic conditions in this period can be primarily attributed to the large number of admissions for diabetes complications. Ear, nose and throat infections; and dehydration and gastroenteritis have the highest rates of avoidable hospitalisations for the acute conditions.

Figure 8: Avoidable hospitalisations¹ by condition, Hawkesbury DGP and New South Wales, 2001/02



¹ Admissions resulting from ACS conditions: excludes nutritional deficiencies as less than ten admissions

Table 7: Avoidable hospitalisations¹ by condition, Hawkesbury DGP, New South Wales and Australia, 2001/02

Sub-category/ condition	Hawkesbury DGP		New South Wales		Australia	
	No.	Rate ²	No.	Rate ²	No.	Rate ²
Vaccine-preventable	51	58.9	5,630	84.5	16,573	85.4
Influenza and pneumonia	37	44.5	4,280	64.1	13,021	67.1
Other vaccine preventable	14	14.4	1,350	20.4	3,552	18.3
Chronic³	1,135	1,599.7	106,803	1,587.0	352,545	1,816
Diabetes complications	389	553.3	34,975	519.5	141,345	728.1
Iron deficiency anaemia	40	54.6	4,494	67.0	16,451	84.7
Hypertension	16	22.2	2,398	35.7	6,354	32.7
Congestive heart failure	124	215.0	14,270	209.7	42,447	218.6
Angina	164	243.0	16,987	251.8	49,963	257.4
Chronic obstructive pulmonary disease	200	316.5	19,359	285.6	54,853	282.6
Asthma	202	195.1	14,289	216.8	41,009	211.3
Acute	880	931.2	62,543	946.0	200,913	1,035
Dehydration and gastroenteritis	125	151.4	11,725	176.4	37,766	194.5
Convulsions and epilepsy	138	139.6	11,093	168.1	31,137	160.4
Ear, nose and throat infections	222	203.9	10,615	161.1	32,075	165.2
Dental conditions	144	139.9	11,196	170.3	43,667	224.9
Perforated/bleeding ulcer	15	21.9	1,830	27.1	5,795	29.9
Ruptured appendix	13	13.6	1,212	18.5	3,866	19.9
Pyelonephritis	25	26.7	2,038	31.0	7,386	38.0
Pelvic inflammatory disease	47	47.0	2,134	32.7	6,547	33.7
Cellulitis	129	155.3	9,451	142.0	28,204	145.3
Gangrene	22	31.9	1,249	18.6	4,470	23.0
Total avoidable hospitalisations⁴	1,981	2,468.4	170,066	2,543.8	552,786	2,847.5

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

³ Excludes nutritional deficiencies as less than ten admissions

⁴ Sub-category and condition numbers and rates do not add to the reported total avoidable admissions: five conditions (influenza & pneumonia, other vaccine preventable, diabetes complications, ruptured appendix and gangrene) are counted in 'any diagnosis', so may be included in more than one condition group

Avoidable mortality

Avoidable and amenable mortality comprises those causes of death that are potentially avoidable at the present time, given available knowledge about social and economic policy impacts, health behaviours, and health care (the latter relating to the subset of amenable causes).

For information on the avoidable and amenable mortality conditions and ICD codes included in the analysis in this section, please refer to the *Australian and New Zealand Atlas of Avoidable Mortality*, available from www.publichealth.gov.au.

Almost three quarters (71.0%) of all deaths in Hawkesbury-Hills DGP at ages 0 to 74 years over the period 1997 to 2001 are considered to be avoidable, marginally lower than the proportion for country New South Wales (71.6%) (Table 8). However, the rate in the Division is notably (17%) lower than that in country New South Wales, a differential of 0.83.

Deaths amenable to health care (amenable mortality, a subset of avoidable mortality) accounted for 30.0% of all deaths at ages 0 to 74 years in Hawkesbury-Hills DGP, compared to 28.3% in country New South Wales.

Table 8: Avoidable and unavoidable mortality (0 to 74 years) by area, Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Mortality category	Hawkesbury-Hills DGP		Country NSW		New South Wales		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable	1,642	195.3	29,442	234.3	66,151	213.6	189,845	211.8
% of total	71.0	..	71.6	..	71.4	..	71.5	..
(Amenable)	(693)	(83.6)	(11,638)	(91.2)	(26,374)	(85.0)	(76,249)	(85.1)
(% of total)	(30.0)	(..)	(28.3)	(..)	(28.5)	(..)	(28.7)	(..)
Unavoidable	671	80.6	11,700	92.1	26,468	85.3	75,582	84.3
% of total	29.0	..	28.4	..	28.6	..	28.5	..
Total mortality	2,312	275.8	41,142	326.4	92,619	299.0	265,427	296.1
%	100.0	..	100.0	..	100.0	..	100.0	..

¹ Rate is the indirectly age-standardised rate per 100,000 population

Rates of avoidable mortality were higher for males than for females in each of the comparator areas. Hawkesbury-Hills DGP's rate of avoidable mortality for males was 248.2 deaths per 100,000 males, notably higher than the rate of 141.7 for females. Similarly, the rate of amenable mortality for males in the Division was higher, 91.7, compared to 75.3 for females, a rate ratio of 1.22 (Figure 9, Table 9).

Figure 9: Avoidable and amenable mortality by sex (0 to 74 years), Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Note: the different scales




Table 9: Avoidable and amenable mortality (0 to 74 years) by sex, Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Mortality category and sex	Hawkesbury-Hills DGP		Country NSW		New South Wales		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
Males	1,063	248.2	19,569	308.5	43,074	276.8	123,026	272.6
Females	578	141.7	9,873	159.1	23,077	149.6	66,819	150.1
Total	1,642	195.3	29,442	234.3	66,151	213.6	189,845	211.8
Rate ratio-M:F²	..	1.75**	..	1.94**	..	1.85**	..	1.82**
Amenable								
Males	384	91.7	6,743	103.6	14,811	94.8	42,568	94.3
Females	309	75.3	4,895	78.6	11,562	74.9	33,681	75.7
Total	693	83.6	11,638	91.2	26,374	85.0	76,249	85.1
Rate ratio-M:F²	..	1.22*	..	1.32**	..	1.27**	..	1.25**

¹ Rate is the indirectly age-standardised rate per 100,000 population

² Rate ratio (M:F) is the ratio of male to female rates; rate ratios differing significantly from 1.0 are shown with * p <0.05; ** p <0.01

Another way of measuring premature mortality is to calculate the number of years of life lost (YLL)¹, which takes into account the years a person could have expected to live at each age of death based on the average life expectancy at that age.

The numbers of YLL for Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia over the period of analysis are shown in Table 10 by mortality category. However, given the substantial variation in the populations of these areas, a comparison of the proportion of YLL for each area is also shown.

YLL from avoidable mortality accounted for 71.1% of total YLL (0 to 74 years) for Hawkesbury-Hills DGP, consistent with the 71.8% for country New South Wales. The proportion of YLL from amenable mortality for Hawkesbury-Hills DGP (29.2%) was higher than that for country New South Wales (27.6%).

Table 10: Years of life lost from avoidable mortality (0 to 74 years), Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Mortality category	Hawkesbury-Hills DGP		Country NSW		New South Wales		Australia	
	No.	% of total	No.	% of total	No.	% of total	No.	% of total
Avoidable	29,959	71.1	502,860	71.8	1,147,183	71.8	3,327,375	71.9
(Amenable)	(12,313)	(29.2)	(192,960)	(27.6)	(444,143)	(27.8)	(1,298,430)	(28.0)
Unavoidable	12,163	28.9	197,182	28.2	451,496	28.2	1,303,289	28.1
Total	42,123	100.0	700,042	100.0	1,598,679	100.0	4,630,664	100.0

¹ Years of life lost were calculated using the remaining life expectancy method (this provides an estimate of the average time a person would have lived had he or she not died prematurely). The reference life table was the Coale and Demeny Model Life Table West level 26 female (for both males and females), with the YLL discounted to net present value at a rate of 3 per cent per year.

In each of the areas in Table 11, the majority of avoidable mortality at ages 0 to 74 years occurred in the 65 to 74 year age group (Table 11), with 1,390.7 deaths per 100,000 population in Hawkesbury-Hills Division. The 45 to 64 year age group accounted for the next highest rate of avoidable death in all of the comparators, with a rate 265.0 in Hawkesbury-Hills Division.

Table 11: Avoidable and amenable mortality by age, Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Mortality category and age (years)	Hawkesbury-Hills DGP		Country NSW		New South Wales		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
0-14	62	25.5	738	29.0	1,836	27.5	5,669	28.8
15-24	88	54.9	938	62.6	2,241	50.9	7,045	52.8
25-44	217	66.4	3,317	99.6	8,119	82.9	24,356	83.9
45-64	612	265.0	9,755	343.5	22,358	311.1	64,282	304.9
65-74	662	1,390.7	14,694	1464.0	31,597	1,375.8	88,493	1,358.1
Total	1,642	195.3	29,442	234.3	66,151	213.6	189,845	211.8
Amenable								
0-24	56	13.6	645	15.5	1,658	14.8	5,083	15.4
25-44	61	18.6	784	23.0	1,878	19.2	5,946	20.5
45-64	284	123.5	4,060	142.9	9,444	131.4	27,464	130.3
65-74	292	615.4	6,148	613.7	13,394	582.9	37,756	579.4
Total	693	83.6	11,638	91.2	26,374	85.0	76,249	85.1

¹ Rate is the indirectly age-standardised rate per 100,000 population

Table 12 shows the number and age-standardised death rate by selected major condition group and selected causes included in the avoidable mortality classification.

The highest rates of avoidable mortality for the selected major condition groups in the Hawkesbury-Hills DGP were for cardiovascular diseases, with a rate of 68.2 deaths per 100,000 population, and, cancer 66.0 deaths per 100,000 population (Table 12, Figure 10). For the selected causes within the condition groups, the two major causes of avoidable mortality were ischaemic heart disease and lung cancer, with rates of 49.2 per 100,000 population and 21.5 per 100,000, respectively.


Table 12: Avoidable mortality (0 to 74 years) by major condition group and selected cause, Hawkesbury-Hills DGP, country New South Wales, New South Wales and Australia, 1997 to 2001

Condition group/ selected cause	Hawkesbury-Hills DGP		Country NSW		New South Wales		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Cancer	541	66.0	9,239	71.9	21,158	68.1	62,338	69.5
Colorectal cancer	116	14.3	1,936	14.9	4,318	13.9	13,008	14.5
Lung cancer	170	21.5	3,314	25.2	7,297	23.4	21,208	23.7
Cardiovascular diseases	530	68.2	10,101	77.0	21,925	70.3	59,945	66.9
Ischaemic heart disease	385	49.2	7,474	57.0	15,935	51.1	43,712	48.8
Cerebrovascular diseases	112	14.7	2,015	15.4	4,656	14.9	12,558	14.0
Respiratory system diseases	89	12.1	2,136	16.0	4,313	13.8	11,612	13.0
Chronic obstructive pulmonary disease	77	10.7	1,966	14.6	3,882	12.4	10,395	11.6
Unintentional injuries	150	14.9	2,027	18.6	4,540	15.0	14,224	15.9
Road traffic injuries	108	10.6	1,279	11.8	2,528	8.4	8,138	9.1
Intentional injuries	110	11.0	1,939	18.1	4,497	14.9	13,891	15.5
Suicide and self inflicted injuries	97	9.7	1,730	16.1	3,941	13.0	12,393	13.8

¹ Rate is the indirectly age-standardised rate per 100,000 population

Rates in the Division were below those in country New South Wales, and below or consistent with those in Australia (Figure 10).

Figure 10: Avoidable mortality (0 to 74 years) by major condition group and selected cause, Hawkesbury-Hills DGP, country New South Wales and Australia, 1997 to 2001



Notes on the data

Data sources and limitations

General

References to 'country New South Wales' relate to New South Wales excluding the Sydney Statistical Division.

Data sources

Table 13 details the data sources for the material presented in this profile.

Table 13: Data sources

Section	Source
Population	
Figures 1 and 2; Table 1	Estimated Resident Population, ABS, 30 June for the periods shown
Figure 3	Estimated Resident Population, ABS, 30 June 2005; Population Projections, ABS, 30 June 2020 (unpublished) ¹
Additional socio-demographic indicators	
Figure 4	ABS SEIFA package, Census 2001
Table 2; Figure 5; Map 1	Jobless families, ABS, 2001 (unpublished)
Table 2; Figure 5; Map 2	Private health insurance, from Hansard
GP services – patient flow/ GP catchment	
Tables 3 and 4	Medicare Australia, 2003/04
Additional prevalence estimates: chronic diseases and risk factors combined	
Figure 6; Table 5	Estimated from 2001 National Health Survey (NHS), ABS (unpublished)
Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions	
Tables 6 and 7; Figures 7 and 8	National Hospital Morbidity Database at Australian Institute of Health & Welfare, 2001/02; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)
Avoidable mortality	
Tables 8, 9, 10, 11 and 12; Figures 9 and 10	ABS Deaths 1997-2001; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)

¹ The projected population at June 2020 is based on the 2002 ERP. As such, it is somewhat dated, and does not take into account more recent demographic trends: it is however the only projection series available at the SLA level for the whole of Australia.

Methods

For background information on the additional prevalence estimates presented in this profile, please refer to the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Please also refer to the November 2005 profile for information on the data converters.

Mapping

In some Divisions the maps may include a very small part of an SLA which has not been allocated any population; or has a population of less than 100 or has less than 1% of the SLAs total population; or there were less than five cases (ie. jobless families, people with health insurance): these areas are mapped with a pattern.

Statistical geography of the Hawkesbury-Hills DGP

For information on the postcodes in the Division, please refer the Department of Health and Ageing website <http://www.health.gov.au/internet/wcms/publishing.nsf/Content/health-pcd-programs-divisions-divspc.htm>; also included in table format in the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Statistical Local Areas (SLAs) are defined by the Australian Bureau of Statistics to produce areas for the presentation and analysis of data. Hawkesbury SLA is mainly comprised within the Division, Blacktown – North and Baulkham Hills are partly within in this Division. Other SLAs which comprise the Division are shown in Table 14.

Table 14: SLAs and population in Hawkesbury-Hills DGP, 2005 on 2001 boundaries

SLA code	SLA name	Per cent of the SLA's population in the Division*	Estimate of the SLA's 2005 population in the Division
10500	Baulkham Hills	58.8	94,784
10751	Blacktown - North	67.6	62,243
<i>13100</i>	<i>Gosford</i>	<i>0.1</i>	<i>141</i>
13800	Hawkesbury	99.4	63,458
14000	Hornsby	2.2	3,516
16250	Parramatta	7.4	11,247
16350	Penrith	2.8	4,933

* Proportions are approximate and are known to be incorrect in some cases, due to errors in the concordance used to allocate CDs to form postal areas. In addition, in a small number of cases, part(s) of an SLA can be allocated to another Division, sometimes several hundred kilometres away. Although adjustments have not been made to the concordance to correct these errors, the affected SLAs are highlighted in the table (shown in bold italic typeface)

Acknowledgements

Funding for these profiles was provided by the Population Health Division of the Department of Health and Ageing (DoHA).

Further developments and updates

When the re-aligned boundaries are released and DoHA have made known their geographic composition, PHIDU will examine the need to revise and re-publish these profiles (*Population health profile*, dated November 2005, and the *Population health profile: supplement*, dated March 2007).

PHIDU contact details

For general comments, data issues or enquiries re information on the web site, please contact PHIDU:

Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au