

Population health profile of the Yorke Peninsula

Division of General Practice: supplement

Population Profile Series: No. 92a

PHIDU

March 2007



PHIDU



Australian Government

Australian Institute of
Health and Welfare



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA



© Commonwealth of Australia 2007

This work may be reproduced and used subject to acknowledgement of the source of any material so reproduced.

National Library of Australia Cataloguing in Publication entry

Population health profile of the Yorke Peninsula Division of General Practice: supplement.

Bibliography.

ISBN 9 78073089 6883 (web).

1. Public health - South Australia - Yorke Peninsula - Statistics. 2. Health status indicators - South Australia - Yorke Peninsula - Statistics. 3. Health service areas - South Australia - Yorke Peninsula. 4. Yorke Peninsula (S. Aust.) - Statistics, Medical. I. Public Health Information Development Unit (Australia). (Series : Population profile series ; no. 92a).

362.10994235

ISSN 1833-0452 Population Profile Series

Public Health Information Development Unit, The University of Adelaide
A Collaborating Unit of the Australian Institute of Health and Welfare

This profile was produced by PHIDU, the Public Health Information Development Unit at The University of Adelaide, South Australia. The work was funded under a grant from the Australian Government Department of Health and Ageing. The views expressed in this profile are solely those of the authors and should not be attributed to the Department of Health and Ageing or the Minister for Health and Ageing.

Interpretation of differences between data in this profile and similar data from other sources needs to be undertaken with care, as such differences may be due to the use of different methodology to produce the data.

Suggested citation:

PHIDU. (2007) *Population health profile of the Yorke Peninsula Division of General Practice: supplement*. Population Profile Series: No. 92a. Public Health Information Development Unit (PHIDU), Adelaide.

Enquiries about or comments on this publication should be addressed to:

PHIDU, The University of Adelaide, South Australia 5005
Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au

This publication, the maps and supporting data, together with other publications on population health, are available from the PHIDU website (www.publichealth.gov.au).

Published by Public Health Information Development Unit, The University of Adelaide

Contributors: Anthea Page, Sarah Ambrose, Kristin Leahy and John Glover

Population health profile

of the Yorke Peninsula Division of General Practice: supplement

This profile is a supplement to the *Population health profile of the Yorke Peninsula Division of General Practice*, dated November 2005, available from www.publichealth.gov.au. This supplement includes an update of the population of the Yorke Peninsula Division of General Practice, as well as additional indicators and aspects of the Division's socioeconomic status, use of GP services and health. The contents are:


- Population [updated to June 2005]
- Additional socio-demographic indicators
- Unreferred attendances – patient flow/ GP catchment
- Additional prevalence estimates: chronic diseases and risk factors combined
- Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions
- Avoidable mortality

For further information on the way Division totals in this report have been estimated, please refer to the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Population

The Yorke Peninsula Division had an Estimated Resident Population of 25,143 at 30 June 2005.

Figure 1: Annual population change, Yorke Peninsula DGP, country South Australia, South Australia and Australia, 1991 to 1996, 1996 to 2001 and 2001 to 2005




Over the five years from 1991 to 1996, the Division's population increased by 0.2% on average each year, consistent with the small increases in country South Australia (0.3%) and South Australia (0.4%). From 1996 to 2001, the annual percentage increase in the Division was 0.4%, equal to that for country South Australia, but less than for South Australia (0.6%). The growth rate of 0.8% per year on average from 2001 to 2005 was above the levels for country South Australia (0.5%) and South Australia (0.4%).

Table 1: Population by age, Yorke Peninsula DGP and Australia, 2005

Age group (years)	Yorke Peninsula DGP		Australia	
	No.	%	No.	%
0-14	4,406	17.5	3,978,221	19.6
15-24	2,259	9.0	2,819,834	13.9
25-44	5,337	21.2	5,878,107	28.9
45-64	7,421	29.5	4,984,446	24.5
65-74	3,089	12.3	1,398,831	6.9
75-84	2,044	8.1	954,143	4.7
85+	587	2.3	315,027	1.5
Total	25,143	100.0	20,328,609	100.0

As shown in the accompanying table and the age-sex pyramid (Figure 2), the Yorke Peninsula DGP had notably fewer young people aged 15 to 24 years (9.0%), and people aged 25 to 44 years (21.2%) than Australia as a whole (with 13.9% and 28.9%) (Table 1). Conversely, the proportions of the Division's population aged 45 years and over were higher than those for Australia.


Figure 2: Population in Yorke Peninsula DGP and Australia, by age and sex, 2005



There are marked differences in the age distribution of the Division's population (when compared to Australia overall). The most notable are:

- at younger ages – fewer children aged 0 to 9 years and young people aged 15 to 19 years;
- from 20 to 49 years – fewer males and females, markedly so between the ages of 20 to 34 years; and
- at older ages – notably higher proportions of males and females at age 50 years and over.

Figure 3: Population projections for Yorke Peninsula DGP, by age and sex, 2005 and 2020




The population projections for the Division show a number of changes in age distribution, with the 2020 population projected to have:

- at younger ages – relatively fewer children and young people aged 0 to 19 years;
- from 20 to 54 years – relatively fewer males and females; and
- from 55 years onwards – relatively more males and females, particularly between 60 and 74 years of age.

Additional socio-demographic indicators

Please refer to the earlier *Population health profile of the Yorke Peninsula Division of General Practice*, dated November 2005, available from www.publichealth.gov.au, for other socio-demographic indicators.

Figure 4: Index of Relative Socio-Economic Disadvantage, Yorke Peninsula DGP, 2001



One of four socioeconomic indexes for areas produced at the 2001 ABS Census is the Index of Relative Socio-Economic Disadvantage.

The Yorke Peninsula DGP has an index score of 987, below the score for Australia of 1000: this score varies across the Division, from a score of 906 in the most disadvantaged areas to 1066 in the least disadvantaged areas.


Note: each 'quintile' comprises approximately 20% of the population of the Division.

A new indicator, produced for the first time at the 2001 ABS Census, shows the number of jobless families with children under 15 years of age. There were notably more jobless families in the Yorke Peninsula DGP (22.7%), compared to country South Australia as a whole (17.5%) (Figure 5, Table 2).

With the introduction of the 30% rebate for private health insurance premiums, there was a once-off registration process, providing information of the postcode and residence of those who had such insurance (these data are not available at this area level for later dates). In 2001, the Division had the same proportion of people with private health insurance (43.5%) as in country South Australia (43.5%) (Figure 5, Table 2).

Figure 5: Socio-demographic indicators, Yorke Peninsula DGP, country South Australia, South Australia and Australia, 2001

Jobless families with children under 15 years old



Private health insurance, 30 June






Table 2: Socio-demographic indicators, Yorke Peninsula DGP, country South Australia, South Australia and Australia, 2001

Indicator	Yorke Peninsula DGP		Country SA		South Australia		Australia	
	No.	%	No.	%	No.	%	No.	%
Jobless families with children under 15 years old	507	22.7	7,725	17.5	29,203	18.7	357,563	17.4
Private health insurance (30 June)	10,239	43.5	173,066	43.5	754,598	51.3	8,671,106	46.0



Details of the distribution of jobless families (Map 1) and of the population covered by private health insurance (Map 2) are shown by Statistical Local Area (SLA) in Maps 1 and 2, respectively.

Map 1: Jobless families with children under 15 years of age by SLA, Yorke Peninsula DGP, 2001



[#] data were not mapped: see 'Mapping' note under Methods

Map 2: People covered by private health insurance by SLA, Yorke Peninsula DGP, 30 June 2001



[#] data were not mapped: see 'Mapping' note under Methods

GP services to residents of the Yorke Peninsula DGP

The following tables include information, purchased from Medicare Australia, of the movement of patients and GPs between Divisions. Note that the data only include unreferred attendances recorded under Medicare: unreferred attendances not included are those for which the cost is met by the Department of Veterans' Affairs or a compensation scheme; or are provided by salaried medical officers in hospitals, community health services or Aboriginal Medical Services, and which are not billed to Medicare. At any attendance, one or more services may have been provided.

The majority (87.9%) of all unreferred attendances to residents of Yorke Peninsula DGP were provided in the Division (ie. by a GP with a provider number in the Division): this represented 118,159 GP unreferred attendances (Table 3). A further 3.8% of unreferred attendances to residents were provided by GPs with a provider number in Mid North Rural DGP.

Table 3: Patient flow – People living¹ in Yorke Peninsula DGP by Division where attendance occurred², 2003/04

Division		Unreferred attendances	
Number	Name	No.	% ³
507	Yorke Peninsula DGP	118,159	87.9
508	Mid North Rural DGP	5,174	3.8
504	Adelaide Central and Eastern DGP	2,152	1.6
502	Adelaide Northern DGP	1,967	1.5
501	Adelaide Western DGP	1,766	1.3
505	Southern DGP	1,613	1.2
503	Adelaide North East DGP	1,199	0.9
Other	..	2,444	1.8
Total	..	134,474	100.0

¹ Based on address in Medicare records

² Division of GP based on provider number

³ Proportion of all unreferred attendances of patients with an address in Division 507 by Division in which attendance occurred

The majority (93.7%) of unreferred attendances provided by GPs with a provider number in Yorke Peninsula DGP were also to people living in the Division (ie. their Medicare address was in the Division) (Table 4). A further 1.1% of unreferred attendances by GPs in the Division were to residents of Southern DGP.

Table 4: GP catchment – Unreferred attendances provided by GPs¹ in Yorke Peninsula DGP by Division of patient address², 2003/04

Division		Unreferred attendances	
Number	Name	No.	% ³
507	Yorke Peninsula DGP	118,159	93.7
505	Southern DGP	1,349	1.1
502	Adelaide Northern DGP	1,042	0.8
503	Adelaide North East DGP	936	0.7
501	Adelaide Western DGP	930	0.7
508	Mid North Rural DGP	844	0.7
504	Adelaide Central and Eastern DGP	703	0.6
Other	..	2180	1.7
Total	..	137,891	100.0

¹ Division of GP based on provider number

² Based on address in Medicare records

³ Proportion of all unreferred attendances to GPs with a provider number in Division 507 by Division of patient address

Additional prevalence estimates: chronic diseases and risk factors combined

Please refer to the earlier *Population health profile of the Yorke Peninsula Division of General Practice*, dated November 2005, available from www.publichealth.gov.au, for the separate prevalence estimates of chronic disease; measures of self-reported health and risk factors. The process by which the estimates have been made, and details of their limitations, are also described in the 'Notes on the data' section of this earlier profile.

In this section two estimates, which combine the prevalence of selected chronic diseases with a risk factor, are shown for the Division. The measures are of people who *had asthma and were smokers*, and people who *had type 2 diabetes and were overweight or obese*: note that the estimates have been predicted from self-reported data, and are not based on clinical records or physical measures.

It is estimated that there were relatively more people in Yorke Peninsula DGP who had asthma and were smokers, compared to country South Australia or Australia as a whole (Figure 6, Table 5): that is, the prevalence rates per 1,000 population were higher. Similarly, there was a higher rate of people in Yorke Peninsula DGP who had type 2 diabetes and were overweight/ obese, compared to country South Australia; however, the rate was lower than for Australia as a whole.

Figure 6: Estimates of selected chronic diseases and risk factors, Yorke Peninsula DGP, country South Australia and Australia, 2001




Table 5: Estimates of selected chronic diseases and risk factors, Yorke Peninsula DGP, country South Australia, South Australia and Australia, 2001

Variable	Yorke Peninsula DGP		Country SA		South Australia		Australia	
	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ²	No. ¹	Rate ¹
Had asthma & smoked ³	542	28.3	9,057	25.5	32,487	22.3	397,734	20.8
Had type 2 diabetes & were overweight/ obese ⁴	479	14.0	5,425	12.8	23,187	14.9	283,176	15.2

¹ No. is a weighted estimate of the number of people in Yorke Peninsula DGP reporting these chronic conditions/ with these risk factors and is derived from synthetic predictions from the 2001 NHS

² Rate is the indirectly age-standardised rate per 1,000 population

³ Population aged 18 years and over

⁴ Population aged 15 years and over

Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions

The rationale underlying the concept of avoidable hospitalisations is that timely and effective care of certain conditions, delivered in a primary care setting, can reduce the risk of hospitalisation. Admissions to hospital for these ambulatory care sensitive (ACS) conditions can be avoided in three ways. Firstly, for conditions that are usually preventable through immunisation or nutritional intervention, disease can be prevented almost entirely. Secondly, diseases or conditions that can lead to rapid onset problems, such as dehydration and gastroenteritis, can be treated. Thirdly, chronic conditions, such as congestive heart failure, can be managed to prevent or reduce the severity of acute flare-ups to avoid hospitalisation.

This measure does not include other aspects of avoidable morbidity, namely potentially preventable hospitalisations (hospitalisations resulting from diseases preventable through population based health promotion strategies, e.g. alcohol-related conditions; and most cases of lung cancer) and hospitalisations avoidable through injury prevention (e.g. road traffic accidents).

For information on the ambulatory care sensitive conditions and ICD codes included in the analysis in this section, please refer to the *Atlas of Avoidable Hospitalisations in Australia: ambulatory care-sensitive conditions*, available from www.publichealth.gov.au.

In 2001 to 2002, the 1,113 admissions from ambulatory care sensitive (ACS) conditions accounted for 11.6% of all admissions in the Yorke Peninsula DGP (Table 6, Figure 7), markedly above the levels in South Australia (8.5) and Australia (8.7%).


Table 6: Avoidable¹ and unavoidable hospitalisations, Yorke Peninsula DGP, South Australia, and Australia, 2001/02

Category	Yorke Peninsula DGP			South Australia			Australia		
	No.	Rate ²	%	No.	Rate ²	%	No.	Rate ²	%
Avoidable ¹	1,113	3,464.9	11.6	47,247	2,915.7	8.5	552,786	2,847.5	8.7
Unavoidable	8,509	28,878.6	88.4	507,053	32,039.4	91.5	5,818,199	29,970.7	91.3
Total	9,622	32,402.5	100.0	554,300	34,952.2	100.0	6,370,985	32,818.2	100.0

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

Figure 7: Avoidable hospitalisations¹, Yorke Peninsula DGP, South Australia and Australia, 2001/02




The rate of avoidable hospitalisations in Yorke Peninsula DGP is markedly higher, a rate of 3,464.9 admissions per 100,000 population, compared to both South Australia (a rate of 2,915.7), Australia (2,847.5).

¹ Admissions resulting from ACS conditions

Diabetes complications; asthma; dehydration and gastroenteritis; ear, nose and throat infections; and chronic obstructive pulmonary disease were the five conditions with the highest rates of avoidable hospitalisations in the Yorke Peninsula DGP (Figure 8, Table 7).

Table 7 shows the number, rate and proportion of avoidable hospitalisations, for the individual ACS conditions, as well as the vaccine-preventable; acute; and chronic sub-categories. The majority of avoidable hospitalisations are attributable to chronic health conditions. The predominance of hospitalisations for chronic conditions in this period can be primarily attributed to the large number of admissions for diabetes complications and asthma. Dehydration and gastroenteritis, and ear, nose and throat infections, have the highest rates of avoidable hospitalisations for the acute conditions.

Figure 8: Avoidable hospitalisations¹ by condition, Yorke Peninsula DGP and South Australia, 2001/02



¹ Admissions resulting from ACS conditions: excludes nutritional deficiencies as less than ten admissions; and other vaccine-preventable conditions and ruptured appendix as number of admissions insufficient

Table 7: Avoidable hospitalisations¹ by condition, Yorke Peninsula DGP, South Australia and Australia, 2001/02

Sub-category/ condition	Yorke Peninsula DGP		South Australia		Australia	
	No.	Rate ²	No.	Rate ²	No.	Rate ²
Vaccine-preventable	28	90.2	1,466	92.9	16,573	85.4
Influenza and pneumonia	28	90.2	1,075	67.0	13,021	67.1
Other vaccine preventable	#	..	391	25.9	3,552	18.3
Chronic³	762	2,211.0	30,607	1,837.6	352,545	1,816
Diabetes complications	227	611.4	11,640	692.9	141,345	728.1
Iron deficiency anaemia	18	53.3	1,271	76.1	16,451	84.7
Hypertension	35	101.6	532	31.6	6,354	32.7
Congestive heart failure	124	315.0	3,900	219.1	42,447	218.6
Angina	96	252.9	3,778	221.6	49,963	257.4
Chronic obstructive pulmonary disease	134	334.1	4,710	272.9	54,853	282.6
Asthma	128	542.7	4,776	323.4	41,009	211.3
Acute	358	1,402.4	16,405	1,077.6	200,913	1,035
Dehydration and gastroenteritis	101	350.8	3,111	194.8	37,766	194.5
Convulsions and epilepsy	38	159.0	2,153	143.6	31,137	160.4
Ear, nose and throat infections	75	345.9	3,046	210.9	32,075	165.2
Dental conditions	66	281.0	3,831	259.2	43,667	224.9
Perforated/bleeding ulcer	8	21.8	555	32.5	5,795	29.9
Ruptured appendix	#	..	255	17.0	3,866	19.9
Pyelonephritis	8	31.3	681	44.7	7,386	38.0
Pelvic inflammatory disease	5	27.1	497	33.7	6,547	33.7
Cellulitis	50	167.2	1,987	124.1	28,204	145.3
Gangrene	7	18.3	289	17.1	4,470	23.0
Total avoidable hospitalisations⁴	1,113	3,464.9	47,247	2,915.7	552,786	2,847.5

¹ Admissions resulting from ACS conditions

² Rate is the indirectly age-standardised rate per 100,000 population

³ Excludes nutritional deficiencies as less than ten admissions

⁴ Sub-category and condition numbers and rates do not add to the reported total avoidable admissions: five conditions (influenza & pneumonia, other vaccine preventable, diabetes complications, ruptured appendix and gangrene) are counted in 'any diagnosis', so may be included in more than one condition group

Not shown or not calculated as there are fewer than five admissions over the period

Avoidable mortality

Avoidable and amenable mortality comprises those causes of death that are potentially avoidable at the present time, given available knowledge about social and economic policy impacts, health behaviours, and health care (the latter relating to the subset of amenable causes).

For information on the avoidable and amenable mortality conditions and ICD codes included in the analysis in this section, please refer to the *Australian and New Zealand Atlas of Avoidable Mortality*, available from www.publichealth.gov.au.

Almost three quarters (74.4%) of all deaths in Yorke Peninsula DGP at ages 0 to 74 years over the period 1997 to 2001 are considered to be avoidable, higher than the proportion for country South Australia (72.5%) (Table 8). Deaths amenable to health care (amenable mortality, a subset of avoidable mortality) accounted for 32.1% of all deaths at ages 0 to 74 years in Yorke Peninsula DGP, compared to 29.8% in country South Australia.

Table 8: Avoidable and unavoidable mortality (0 to 74 years) by area, Yorke Peninsula DGP, country South Australia, South Australia and Australia, 1997 to 2001

Mortality category	Yorke Peninsula DGP		Country SA		South Australia		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable	413	246.8	4,852	230.3	15,938	210.4	189,845	211.8
% of total	74.4	..	72.5	..	71.4	..	71.5	..
(Amenable)	(178)	(103.5)	(1,993)	(93.6)	(6,556)	(85.9)	(76,249)	(85.1)
(% of total)	(32.1)	(..)	(29.8)	(..)	(29.4)	(..)	(28.7)	(..)
Unavoidable	142	83.3	1,837	86.5	6,369	83.7	75,582	84.3
% of total	25.6	..	27.5	..	28.6	..	28.5	..
Total mortality	555	329.8	6,688	316.8	22,307	294.1	265,427	296.1
%	100.0	..	100.0	..	100.0	..	100.0	..

¹ Rate is the indirectly age-standardised rate per 100,000 population

Rates of avoidable mortality were higher for males than for females in each of the comparator areas. Yorke Peninsula DGP's rate of avoidable mortality for males was 319.2 deaths per 100,000 males, higher than the rate of 173.2 for females. Similarly, the rate of amenable mortality for males in the Division was higher, 114.6, compared to 92.2 for females, a rate ratio of 1.24 (Figure 9, Table 9).

Figure 9: Avoidable and amenable mortality by sex (0 to 74 years), Yorke Peninsula DGP, country South Australia, South Australia and Australia, 1997 to 2001

Note: the different scales




Table 9: Avoidable and amenable mortality (0 to 74 years) by sex, Yorke Peninsula DGP, country South Australia, South Australia and Australia, 1997 to 2001

Mortality category and sex	Yorke Peninsula DGP		Country SA		South Australia		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
Males	276	319.2	3,259	300.9	10,326	272.8	123,026	272.6
Females	137	173.2	1,593	158.3	5,612	147.2	66,819	150.1
Total	413	246.8	4,852	230.3	15,938	210.4	189,845	211.8
Rate ratio-M:F²	..	1.84**	..	1.90**	..	1.85**	..	1.82**
Amenable								
Males	105	114.6	1,169	105.6	3,671	96.0	42,568	94.3
Females	73	92.2	824	81.4	2,884	75.7	33,681	75.7
Total	178	103.5	1,993	93.6	6,556	85.9	76,249	85.1
Rate ratio-M:F²	..	1.24	..	1.30**	..	1.27**	..	1.25**

¹ Rate is the indirectly age-standardised rate per 100,000 population

² Rate ratio (M:F) is the ratio of male to female rates; rate ratios differing significantly from 1.0 are shown with * p <0.05; ** p <0.01

Another way of measuring premature mortality is to calculate the number of years of life lost (YLL)¹, which takes into account the years a person could have expected to live at each age of death based on the average life expectancy at that age.

The numbers of YLL for Yorke Peninsula DGP, country South Australia, South Australia and Australia over the period of analysis are shown in Table 10 by mortality category. However, given the substantial variation in the populations of these areas, a comparison of the proportion of YLL for each area is also shown.

YLL from avoidable mortality accounted for 74.3% of total YLL (0 to 74 years) for Yorke Peninsula DGP, higher than the 72.9% for country South Australia. The proportion of YLL from amenable mortality of 31.8% for Yorke Peninsula DGP was higher than the 28.9% for country South Australia.

Table 10: Years of life lost from avoidable mortality (0 to 74 years), Yorke Peninsula DGP, country South Australia, South Australia and Australia, 1997 to 2001

Mortality category	Yorke Peninsula DGP		Country SA		South Australia		Australia	
	No.	% of total	No.	% of total	No.	% of total	No.	% of total
Avoidable	6,570	74.3	83,705	72.9	273,135	71.8	3,327,375	71.9
(Amenable)	(2,808)	(31.8)	(33,165)	(28.9)	(108,777)	(28.6)	(1,298,430)	(28.0)
Unavoidable	2,269	25.7	31,059	27.1	107,223	28.2	1,303,289	28.1
Total	8,839	100.0	114,764	100.0	380,358	100.0	4,630,664	100.0

¹ Years of life lost were calculated using the remaining life expectancy method (this provides an estimate of the average time a person would have lived had he or she not died prematurely). The reference life table was the Coale and Demeny Model Life Table West level 26 female (for both males and females), with the YLL discounted to net present value at a rate of 3 per cent per year.

In each of the areas in Table 11, the majority of avoidable mortality at ages 0 to 74 years occurred in the 65 to 74 year age group (Table 11), with 1,519.5 deaths per 100,000 population in the Yorke Peninsula Division. The 45 to 64 year age group accounted for the next highest rate of avoidable death in all of the comparators, with a rate of 384.6 in the Yorke Peninsula Division.

Table 11: Avoidable and amenable mortality by age, Yorke Peninsula DGP, country South Australia, South Australia and Australia, 1997 to 2001

Mortality category and age (years)	Yorke Peninsula DGP		Country SA		South Australia		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Avoidable								
0-14	7	34.8	118	26.8	352	24.2	5,669	28.8
15-24	4	42.3	159	67.5	523	52.4	7,045	52.8
25-44	29	99.7	596	99.3	1,979	88.8	24,356	83.9
45-64	141	384.6	1,640	333.3	5,130	297.8	64,282	304.9
65-74	231	1,519.5	2,338	1439.0	7,954	1354.8	88,493	1,358.1
Total	413	246.8	4,852	230.3	15,938	210.4	189,845	211.8
Amenable								
0-24	6	16.4	101	14.1	324	13.3	5,083	15.4
25-44	9	29.0	146	23.8	507	22.6	5,946	20.5
45-64	62	168.1	710	144.8	2,248	130.1	27,464	130.3
65-74	102	678.6	1,036	641.3	3,477	591.6	37,756	579.4
Total	178	103.5	1,993	93.6	6,556	85.9	76,249	85.1

¹ Rate is the indirectly age-standardised rate per 100,000 population

Table 12 shows the number and age-standardised death rate by selected major condition group and selected causes included in the avoidable mortality classification.

The highest rates of avoidable mortality for the selected major condition groups in the Yorke Peninsula DGP were for cardiovascular diseases, a rate of 87.2 deaths per 100,000 population, and cancer, with a rate of 73.7 deaths per 100,000 population (Table 12, Figure 10). For the selected causes within the condition groups, the two major causes of avoidable mortality were ischaemic heart disease and lung cancer, with rates of 67.4 per 100,000 population and 24.3 per 100,000, respectively.


Table 12: Avoidable mortality (0 to 74 years) by major condition group and selected cause, Yorke Peninsula DGP, country South Australia, South Australia and Australia, 1997 to 2001

Condition group/ selected cause	Yorke Peninsula DGP		Country SA		South Australia		Australia	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Cancer	132	73.7	1,489	69.4	5,209	67.8	62,338	69.5
Colorectal cancer	31	17.2	346	16.0	1,142	14.8	13,008	14.5
Lung cancer	46	24.3	477	21.8	1,728	22.3	21,208	23.7
Cardiovascular diseases	163	87.2	1,669	76.5	5,324	68.5	59,945	66.9
Ischaemic heart disease	126	67.4	1,260	57.6	3,918	50.5	43,712	48.8
Cerebrovascular diseases	30	16.0	316	14.6	1,086	13.9	12,558	14.0
Respiratory system diseases	30	15.6	270	12.3	897	11.4	11,612	13.0
Chronic obstructive pulmonary disease	25	12.5	239	10.8	783	9.9	10,395	11.6
Unintentional injuries	24	22.5	412	22.1	1,085	15.5	14,224	15.9
Road traffic injuries	17	16.3	307	16.6	687	9.9	8,138	9.1
Intentional injuries	7	7.1	329	17.5	1,138	16.3	13,891	15.5
Suicide and self inflicted injuries	7	6.9	293	15.6	1,018	14.5	12,393	13.8

¹ Rate is the indirectly age-standardised rate per 100,000 population

Rates in the Division were above those for country South Australia and Australia for all condition groups and selected causes other than intentional injuries (Figure 10).

Figure 10: Avoidable mortality (0 to 74 years) by major condition group and selected cause, Yorke Peninsula DGP, country South Australia and Australia, 1997 to 2001



Notes on the data

Data sources and limitations

General

References to 'country South Australia' relate to South Australia excluding the Adelaide Statistical Division.

Data sources

Table 13 details the data sources for the material presented in this profile.

Table 13: Data sources

Section	Source
Population	
Figures 1 and 2; Table 1	Estimated Resident Population, ABS, 30 June for the periods shown
Figure 3	Estimated Resident Population, ABS, 30 June 2005; Population Projections, ABS, 30 June 2020 (unpublished) ¹
Additional socio-demographic indicators	
Figure 4	ABS SEIFA package, Census 2001
Table 2; Figure 5; Map 1	Jobless families, ABS, 2001 (unpublished)
Table 2; Figure 5; Map 2	Private health insurance, from Hansard
GP services – patient flow/ GP catchment	
Tables 3 and 4	Medicare Australia, 2003/04
Additional prevalence estimates: chronic diseases and risk factors combined	
Figure 6; Table 5	Estimated from 2001 National Health Survey (NHS), ABS (unpublished)
Avoidable hospitalisations: hospital admissions resulting from ambulatory care sensitive conditions	
Tables 6 and 7; Figures 7 and 8	National Hospital Morbidity Database at Australian Institute of Health & Welfare, 2001/02; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)
Avoidable mortality	
Tables 8, 9, 10, 11 and 12; Figures 9 and 10	ABS Deaths 1997-2001; data produced in HealthWIZ by Prometheus Information (not available in public release dataset)

¹ The projected population at June 2020 is based on the 2002 ERP. As such, it is somewhat dated, and does not take into account more recent demographic trends: it is however the only projection series available at the SLA level for the whole of Australia.

Methods

For background information on the additional prevalence estimates presented in this profile, please refer to the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Please also refer to the November 2005 profile for information on the data converters.

Mapping

In some Divisions the maps may include a very small part of an SLA which has not been allocated any population; or has a population of less than 100 or has less than 1% of the SLAs total population; or there were less than five cases (i.e. jobless families, people with health insurance): these areas are mapped with a pattern.

Statistical geography of the Yorke Peninsula DGP

For information on the postcodes in the Division, please refer the Department of Health and Ageing website <http://www.health.gov.au/internet/wcms/publishing.nsf/Content/health-pcd-programs-divisions-divspc.htm>; also included in table format in the 'Notes on the data' section of the *Population health profile*, November 2005 (www.publichealth.gov.au).

Statistical Local Areas (SLAs) are defined by the Australian Bureau of Statistics to produce areas for the presentation and analysis of data. In this Division, Yorke Peninsula Local Government Area (LGA) is split into two SLAs – North and South, wholly included in the Division; and a small part of one of the two Port Pirie SLAs is also in the Division. These SLAs and all or parts of the other SLAs listed in Table 14 comprise the Division.

Table 14: SLAs and population in Yorke Peninsula DGP, 2005 on 2001 boundaries

SLA code	SLA name	Per cent of the SLA's population in the Division*	Estimate of the SLA's 2005 population in the Division
40430	Barunga West	41.6	1,072
41560	Copper Coast	100.0	11,640
48130	Wakefield	10.8	711
48831	Yorke Peninsula - North	100.0	7,606
48834	Yorke Peninsula - South	100.0	4,114

* Proportions are approximate and are known to be incorrect in some cases, due to errors in the concordance used to allocate CDs to form postal areas

Acknowledgements

Funding for these profiles was provided by the Population Health Division of the Department of Health and Ageing (DoHA).

Further developments and updates

When the re-aligned boundaries are released and DoHA have made known their geographic composition, PHIDU will examine the need to revise and re-publish these profiles (*Population health profile*, dated November 2005, and the *Population health profile: supplement*, dated March 2007).

PHIDU contact details

For general comments, data issues or enquiries re information on the web site, please contact PHIDU:

Phone: 08-8303 6236 or e-mail: PHIDU@publichealth.gov.au